

Finite Element based Level Set Methods for Higher Order Flows

Martin Burger ^{*} Christina Stöcker [†] Axel Voigt [‡]

March 15, 2007

Abstract

In this paper we shall discuss the numerical simulation of higher order geometric flows by level set methods. Main examples under considerations are surface diffusion and the Willmore flow as well as variants and of them with more complicated surface energies. Such problems find various applications, e.g. in materials science (crystal growth, thin film technology), biophysics (membrane shapes), and computer graphics (surface smoothing and restoration)

We shall use spatial discretizations by finite element methods and semi-implicit time stepping based on local variational principles, which allows to maintain dissipation properties of the flows by the discretization. In order to compensate for the missing maximum principle, which is indeed a major hurdle for the application of level set methods to higher order flows, we employ frequent redistancing of the level set function. We shall review suitable schemes used for redistancing in two and three spatial dimensions.

Finally we also discuss the solution of the arising discretized linear systems in each time step and some particular advantages of the finite element approach such as the possibility of local adaptivity around the zero level set.

Keywords: Level Set Methods, Higher-order Geometric Flows, Finite Element Methods, Semi-Implicit Time Stepping, Energy Dissipation.

^{*}Institut für Numerische und Angewandte Mathematik, Westfälische Wilhelms Universität (WWU), Einsteinstr. 62, D 48149 Münster, Germany. e-mail: martin.burger@uni-muenster.de

[†]Crystal Growth group, Research Center caesar, Ludwig-Erhard-Allee 2, 53175 Bonn, Germany. e-mail: stoecker@caesar.de

[‡]Crystal Growth group, Research Center caesar, Ludwig-Erhard-Allee 2, 53175 Bonn, Germany and Institut für Wissenschaftliche Rechnen, Technische Universität Dresden, Zellescher Weg 12-14, 01062 Dresden, Germany. e-mail: voigt@caesar.de

1 Introduction

Higher order geometric flows received growing attention in the recent years due to emerging applications in various fields, where higher order typically means higher than second order intrinsic derivatives of the surface appearing in the evolution equations. Significant advances have been made in the modelling and understanding of certain flows, while the rigorous mathematical analysis and the construction of flexible computational schemes still lack behind. In particular the use of level set methods, which became popular for lower order flows in the last decade, is not yet well understood. Major obstacles for level set methods in higher order flows are the absence of maximum and comparison principles, which excludes global level set approaches, and the stiffness of the problems, which excludes explicit time stepping to a large extent.

In this paper we develop level set methods for the simulation of higher-order flows based on finite element discretizations in space and semi-implicit time discretizations. The issue of the missing maximum principle is resolved by frequent redistancing. This corresponds to a paradigm for local level set formulations in the continuous setting: a level set solution to a higher-order flow would be an evolving signed distance function that satisfies the geometric evolution equation on the interface (the zero level set) only.

The basis of our approach is a variational formulation of the geometric flows as metric gradient flows (cf. [1, 2, 14]). This means that the computation of an evolving (embedded) curve or surface $\Gamma(t)$ is determined by an energy E (defined on a class of shapes) and a metric structure on a manifold of curves and surfaces. The evolution is determined via the normal velocity v_n , which satisfies an equation of the form

$$\langle v_n, \psi \rangle_{M(\Gamma(t))} = -e'[\Gamma(t)]\psi, \quad (1)$$

for all suitable test velocities ψ , where the scalar product is determined by the metric structure $M(\Gamma(t))$. The right-hand side $e'[\Gamma(t)]\psi$ denotes the variation of the energy functionals with respect to infinitesimal perturbations of the shape with a vector field ψn (which can also be interpreted as a shape derivative, cf. [22]), where n denotes the unit outward normal. Under suitable conditions the shape derivative $e'[\Gamma(t)]$ is a continuous linear functional on the space of test velocities. The most important choices of scalar product in this context are the L^2 and H^{-1} scalar product. In the case of $M(\Gamma)$ denoting the L^2 -scalar product on $\Gamma(t)$, we obtain

$$\int_{\Gamma(t)} v_n \psi \, d\sigma = -e'[\Gamma(t)] \quad \forall \psi \in L^2(\Gamma(t)).$$

Hence, the normal velocity is just the L^2 -representation of the linear functional $e'[\Gamma(t)]$. In the case of e being the area functional, i.e.

$$e[\Gamma] = \int_{\Gamma} 1 \, d\sigma,$$

the variation is computed via the mean curvature h of the surface as

$$e'[\Gamma]\psi = \int_{\Gamma} h\psi \, d\sigma.$$

Thus, one sees that the associated L^2 -gradient flow yields the mean curvature flow $v_n = -h$.

In the case of the H^{-1} -scalar product we have

$$\langle v_n, \psi \rangle_{M(\Gamma(t))} = \int_{\Gamma(t)} \nabla_{\sigma} W_{v_n} \cdot \nabla_{\sigma} W_{\psi} \, d\sigma, \quad (2)$$

with the tangential gradient ∇_{σ} and W_{ψ} defined by the Laplace-Beltrami equation

$$-\nabla_{\sigma} \cdot (\nabla_{\sigma} W_{\psi}) = \psi, \quad \text{on } \Gamma(t) \quad (3)$$

with the additional normalization condition $\int_{\Gamma(t)} W_{\psi} \, d\sigma = 0$. In the case of the area functional the equation for the normal velocity becomes

$$\begin{aligned} - \int_{\Gamma(t)} W_{v_n} \psi \, d\sigma &= - \int_{\Gamma(t)} W_{v_n} \nabla_{\sigma} \cdot (\nabla_{\sigma} \psi) \, d\sigma = \int_{\Gamma(t)} \nabla_{\sigma} W_{v_n} \cdot \nabla_{\sigma} W_{\psi} \, d\sigma \\ &= \int_{\Gamma(t)} h\psi \, d\sigma \end{aligned}$$

and hence, $W_{v_n} = -h$. An application of the Laplace-Beltrami operator finally yields $v_n = \nabla_{\sigma} \cdot (\nabla_{\sigma} h)$, i.e. motion by the Laplacian of curvature, which is the simplest form of a surface diffusion law. Thus, we see that the choice of the H^{-1} scalar product automatically leads to higher order flows, in the above example to a fourth order one.

Another source for higher order geometric flows are simply higher order terms in the surface energy, such as curvatures or even intrinsic derivatives of curvatures. A simple and well-known example is the so-called *Willmore energy*

$$e[\Gamma] = \frac{1}{2} \int_{\Gamma} h^2 \, d\sigma. \quad (4)$$

Since the energy includes second order derivatives via the mean curvature h , its variation (more precisely the L^2 representation) corresponds to a fourth

order differential operator. Thus, in the L^2 scalar product one obtains a fourth order flow, the so-called *Willmore flow*, and by analogy to the above computation we even obtain a sixth order flow in the H^{-1} scalar product.

The paper is organized as follows: In Section 2 we derive level set formulations for the geometric evolution laws, directly from the metric gradient flow formulation. In Section 3 we introduce the finite element formulation for spatial discretization, which is indeed straight-forward. As a consequence of the gradient flow formulation, we construct local-in-time variational problems corresponding to implicit time discretizations, and by further quadratic approximations of the energy functional we obtain semi-implicit schemes in Section 4. In Section 5 we discuss some aspects of redistancing on triangular finite element grids in two and three spatial dimensions. Details about implementation and numerical results are presented in Section 6.

2 Level set formulation of the models

In this section we discuss the level set formulation of the higher order flows distinguishing two cases, namely lower order energies such as the area functional or classical surface energies appearing in materials science, and higher order energies such as the Willmore energy. This distinction is due to the fact that we try to split the higher-order flows into second order systems by the introduction of new variables such as the mean curvature (and constraints defining the new variables in terms of the level set function). In the case of low order energies these variables have to be introduced only in order to approximate the metric, which happens in a very natural way, whereas in the case of higher order energies additional variables are needed in dependence on the metric. In both cases, the constraints can be modeled by using associated Lagrange functionals, the Lagrangian variables will turn out to obey very simple relations in several cases and sometimes also provide interesting new quantities as we shall see below.

2.1 Basic Notations and Level Set Formulations

We start with some basic notations for level set formulations, avoiding a too detailed discussion of level set methods at this point (cf. [37, 36] for further detail). We assume $\Gamma(t) \subset \Omega \subset \mathbb{R}^d$ to be the union of evolving curves or surfaces, which are embedded, i.e.

$$\Gamma(t) = \partial\Theta(t) \setminus \partial\Omega.$$

The level set approach chooses a function $\phi : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\Gamma(t) = \{\phi(\cdot, t) = 0\}, \quad \Theta(t) = \{\phi(\cdot, t) < 0\}.$$

It is well-known (cf. e.g. [36]) that the unit outward normal vector n (pointing into $\Omega \setminus \Theta(t)$), the normal velocity v , and the mean curvature h of the evolving surface can be represented as

$$n = \frac{\nabla\phi}{|\nabla\phi|}|_{\Gamma(t)}, \quad v_n = -\frac{\partial_t\phi}{|\nabla\phi|}|_{\Gamma(t)}, \quad h = \nabla \cdot \left(\frac{\nabla\phi}{|\nabla\phi|} \right)|_{\Gamma(t)}, \quad (5)$$

where we assume that $\nabla\phi \neq 0$. Note that with this sign convention a closed sphere with outer normal n has positive mean curvature $h > 0$. For simplicity we impose in the following natural boundary conditions for ϕ (and other variables) on the boundary of the computational domain $\partial\Omega$.

2.2 Level Set Formulation I: Low Order Energies

We start with low order energies, i.e., energy functionals of the form

$$e[\Gamma] = \int_{\Theta} f(x) + \int_{\Gamma} g(x, n) \, d\sigma, \quad (6)$$

where n denotes the outward unit normal. In many cases f and also parts of g arise through nonlocal interactions, usually modelled via partial differential equations with boundaries or coefficients dependent on Γ . However, such situations can be incorporated via Lagrange functionals, so that the effective functional to be minimized or differentiated with respect to the shape variable always ends up in the form (6).

In the level set context we average the energy over all level sets of ϕ , which yields a new energy on all level sets, respectively on the level set function ϕ directly. For the sake of simplicity we assume that extensions $F : \Omega \rightarrow \mathbb{R}$ and $G : \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}$ of f and g are known (with G being one-homogeneous in the second variable), if not the extensions can be computed in a rather standard way (e.g. constant in normal direction, cf. [36]). Then we obtain the energy from the layer-cake and co-area formula as

$$\begin{aligned} E[\phi] &= \int_{-\infty}^{\phi_{\max}} \int_{\{\phi=\alpha\} \cap \Omega} G(x, \frac{\nabla\phi}{|\nabla\phi|}) \, d\sigma \, d\alpha + \int_{-\infty}^{\phi_{\max}} \int_{\{\phi<\alpha\} \cap \Omega} F(x) \, dx \, d\alpha \\ &= \int_{\Omega} G(x, \nabla\phi) \, dx + \int_{\Omega} F(x)(\phi_{\max} - \phi(x)) \, dx, \end{aligned}$$

where ϕ_{\max} is greater than the maximal value of ϕ appearing in Ω . Since constant terms do not matter for the minimization and variations with respect to ϕ we define the averaged energy as

$$E[\phi] = \int_{\Omega} G(x, \nabla\phi) dx - \int_{\Omega} F(x)\phi(x) dx. \quad (7)$$

In order to compute variations in normal directions we recall the basic level set equation

$$\partial_t\phi = -V_n|\nabla\phi|,$$

for V_n being an extension of the normal velocity v_n . Hence, we compute variations of ϕ only for directions of the form $-V_n|\nabla\phi|$. It is easy that the identity

$$E'[\phi](-V_n|\nabla\phi|) = \int_{-\infty}^{\phi_{\max}} e'[\{\phi = \alpha\}]V_n d\alpha \quad (8)$$

holds for such variations.

In a similar way we can average the scalar products used in the definitions of the geometric flows. The L^2 scalar product can be averaged with the co-area formula as

$$\int_{-\infty}^{\phi_{\max}} \int_{\{\phi=\alpha\}} \psi_1\psi_2 d\sigma d\alpha = \int_{\Omega} \psi_1\psi_2|\nabla\phi| dx = \langle\psi_1, \psi_2\rangle_{\phi}. \quad (9)$$

For the H^{-1} scalar product we use the weak formulation of the Laplace Beltrami Operator and the simple relation

$$\nabla_{\sigma}W = \mathbf{P}\nabla W, \quad \mathbf{P} = I - n \otimes n = I - \frac{\nabla\phi}{|\nabla\phi|} \otimes \frac{\nabla\phi}{|\nabla\phi|}$$

for the tangential derivative to obtain the scalar product $\langle\psi_1, \psi_2\rangle_{\phi}$ as

$$\int_{-\infty}^{\phi_{\max}} \int_{\{\phi=\alpha\}} \nabla_{\sigma}W_{\psi_1} \nabla_{\sigma}W_{\psi_2} d\sigma d\alpha = \int_{\Omega} (\mathbf{P}\nabla W_{\psi_1}) \cdot (\mathbf{P}\nabla W_{\psi_2})|\nabla\phi| dx, \quad (10)$$

where a function W_{ψ} is defined as a weak solution of the (averaged) Laplace-Beltrami equation, i.e.

$$\int_{\Omega} (\mathbf{P}\nabla W_{\psi}) \cdot (\mathbf{P}\nabla\eta)|\nabla\phi| dx = \int_{\Omega} \psi\eta|\nabla\phi| dx \quad (11)$$

for all suitable test functions η .

Now we can simply reformulate the geometric gradient flow as a metric gradient flow on the level set functions via (note that for a variation ψ , the function $-\frac{\psi}{|\nabla\phi|}$ is an extended normal variation)

$$\langle V_n, -\frac{\psi}{|\nabla\phi|} \rangle_\phi = -\langle \frac{\partial_t\phi}{|\nabla\phi|}, -\frac{\psi}{|\nabla\phi|} \rangle_\phi = -E'[\phi](\psi),$$

or, equivalently

$$\langle \frac{\partial_t\phi}{|\nabla\phi|}, \frac{\psi}{|\nabla\phi|} \rangle_\phi = -E'[\phi](\psi). \quad (12)$$

In the case of the area functional ($F \equiv 0$, $G \equiv 1$) and the L^2 scalar product we thus obtain the well-known (weak) level set formulation of the mean curvature flow

$$\int_\Omega \frac{\partial_t\phi\psi}{|\nabla\phi|} dx = - \int_\Omega \frac{\nabla\phi \cdot \nabla\psi}{|\nabla\phi|} dx = \int_\Omega H\psi dx, \quad (13)$$

for all suitable test functions ψ , where $H = \nabla \cdot (\frac{\nabla\phi}{|\nabla\phi|})$ is the extended mean curvature. For the H^{-1} scalar product we use the variable $W := W_{V_n}$ and the identity

$$\int_\Omega (\mathbf{P}\nabla W) \cdot (\mathbf{P}\nabla W - \frac{\psi}{|\nabla\phi|} |\nabla\phi|) dx = - \int_\Omega W\psi dx$$

to obtain in an analogous way the (weak) level set formulation of the simple surface diffusion law, i.e.,

$$\int_\Omega W\psi dx = \int_\Omega \frac{\nabla\phi \cdot \nabla\psi}{|\nabla\phi|} dx \quad (14)$$

$$\int_\Omega (\mathbf{P}\nabla W) \cdot (\mathbf{P}\nabla\eta) |\nabla\phi| dx = - \int_\Omega \partial_t\phi\eta dx \quad (15)$$

for suitable test functions ψ and η . One observes that we have automatically splitted the problem in a system of second order equations incorporating the new variable W , which actually equals the extended mean curvature H .

We mention that the derivation of the level set formulations could also be made by directly rewriting the evolution law in terms of the level set function. For complicated energies, this procedure is rather tedious since one needs to compute the shape derivative first (which can be a hard task for itself) and then one has to compute transformations of higher derivatives to the level set form. This can be avoided in the above approach, one

can directly convert the energy to level set form and then compute classical variations in a function setting (with particular form of the variations). Clearly the above derivation also provides a natural weak level set formulation, which directly leads to finite element discretizations of the equations. Also these formulations have been derived by sometimes tedious calculations in previous work (cf. e.g [19, 24]).

For more general metrics we can formulate the above procedure as rewriting the metric via a Lagrange-functional of the form

$$\left\langle \frac{\partial_t \phi}{|\nabla \phi|}, \frac{\psi}{|\nabla \phi|} \right\rangle_\phi = \partial_\phi L[\partial_t \phi, W, Q; \phi] \psi, \quad (16)$$

where W is the collection of $m_W \in \mathbb{N}_0$ new variables introduced to rewrite the metric and P are the m_W Lagrange multipliers (dual variables) associated to the new constraints defining W . These variables are defined via the conditions

$$\partial_W L[\partial_t \phi, W, Q; \phi] = 0, \quad \partial_Q L[\partial_t \phi, W, Q; \phi] = 0.$$

In the case of the H^{-1} scalar product above we have

$$L[\psi, W, Q; \phi] = \frac{1}{2} \int_\Omega |\mathbf{P}\nabla W|^2 dx + \int_\Omega ((\mathbf{P}\nabla W)(\mathbf{P}\nabla Q) + \psi Q) dx.$$

It is easy to see from the specific form of the functional that the condition $\partial_W L[\partial_t \phi, W, Q; \phi] = 0$ implies $Q = -W$. Moreover, $\partial_Q L[\partial_t \phi, W, Q; \phi] = 0$ yields (15). Finally, inserting $Q = -W$ in (16) and (12) yields (14).

We shall assume in the following that L depends at most quadratically on $\partial_t \phi$ and W and linearly on Q . The assumptions on ∂_t and W are easy to achieve due to the bilinear nature of the scalar product, and the linearity with respect to the dual variable Q is a natural property of Lagrange functionals. The only nonlinear dependence in L thus appears with respect to the level set function ϕ itself, which can be taken care of easily in the time discretization.

2.3 Level Set Formulation II: Higher Order Energies

The second case we consider are higher-order energies such as the Willmore functional. Then we have to introduce $m_U \in \mathbb{N}_0$ further variables U for the splitting and another Lagrangian functional Λ in order to take care of the constraints defining the new variables (and m_U associated Lagrangian or dual variables R). Hence, we shall write

$$E[\phi] = \Lambda[\phi, U, R], \quad (17)$$

with U and R being defined via

$$\partial_U \Lambda[\phi, U, R] = 0, \quad \partial_R \Lambda[\phi, U, R] = 0. \quad (18)$$

Consequently, with the scalar product defined through a Lagrange functional L we equivalently rewrite (12) as

$$\partial_\psi L[\partial_t \phi, W, Q; \phi] = -\partial_\phi \Lambda(\phi, W, Q). \quad (19)$$

We illustrate the procedure for the L^2 gradient flow of the (level set averaged) Willmore functional

$$E[\phi] = \frac{1}{2} \int_{\Omega} H^2 |\nabla \phi| \, dx = \int_{-\infty}^{\phi_{\max}} \int_{\{\phi=\alpha\}} h^2 \, d\sigma \, d\alpha.$$

Here we can simply introduce the variable $U = H$ and the Lagrange functional

$$\Lambda[\phi, U, R] = \frac{1}{2} \int_{\Omega} U^2 |\nabla \phi| \, dx + \int_{\Omega} \left(-UR + \frac{\nabla \phi \cdot \nabla R}{|\nabla \phi|} \right) \, dx.$$

Note that $\partial_R \Lambda[\phi, U, R] = 0$ yields exactly the weak formulation of the equation $U = \nabla \cdot \left(\frac{\nabla \phi}{|\nabla \phi|} \right)$, which defines mean curvature. Moreover, from $\partial_U \Lambda[\phi, U, R] = 0$ we conclude $R = U|\nabla \phi|$. Thus, the dual variable is exactly the *curvature concentration* introduced for the level set formulation of the Willmore flow directly in [24]. One could now use this identity to eliminate U in favour of R as in [24] and one observes that in the general approach above it does not matter which variable we choose for splitting. If we would choose $U = H|\nabla \phi|$ directly as the curvature concentration, then we would obtain the mean curvature as the dual variable.

2.4 Energy Dissipation for the Level Set Formulation

In the following we verify the energy dissipation for (19). For this sake use the test functions $\partial_t \phi$, then we have

$$\begin{aligned} \langle \partial_t \phi, \partial_t \phi \rangle_{\phi} &= \partial_\psi L[\partial_t \phi, W, Q; \phi] \partial_t \phi \\ &= -\partial_\phi \Lambda[\phi, U, R] \partial_t \phi \\ &= -E'[\phi] \partial_t \phi = -\frac{d}{dt} E[\Phi], \end{aligned}$$

or in other words,

$$\frac{d}{dt} E[\Phi] = -\langle \partial_t \phi, \partial_t \phi \rangle_{\phi} = -\|\partial_t \phi\|_{\phi}^2. \quad (20)$$

The energy dissipation property will play a central role in the later time discretization of the flows.

3 Spatial Discretization

In the following we discuss the spatial discretization of the higher order flows or more precisely their level set formulation (19) introduced above. We assume that the new splitting variables have been introduced such that the highest arising differential operators are of second order, which means that only up to first derivatives of all variables appear in the Lagrange functionals L and Λ . Consequently, and also due to the dissipative nature of the flows, it is natural to use finite element methods (with standard Lagrangian elements for time discretization) for the spatial semi-discretization, which we shall briefly discuss in this section.

We choose appropriate finite element subspaces $\mathcal{V}^h \subset \mathcal{V}$ and look for weak solutions satisfying the variational problems in a product of this subspace. Here we shall use $\mathcal{V} = H^1(\Omega)$ and the standard piecewise linear elements

$$\mathcal{V}^h = \{ v \in C(\Omega) \mid v|_T \text{ is linear polynomial for } T \in \mathcal{T} \}, \quad (21)$$

where \mathcal{T} is a decomposition of the polygonal domain Ω into triangles (2D) or tetrahedra (3D). Moreover we assume natural boundary conditions for all variables on the boundary $\partial\Omega$. For the sake of simplicity we assume here that all variables can be discretized robustly by the same kind of elements, i.e., we look for

$$(\phi^h, W^h, U^h, Q^h, R^h) \in \mathcal{P}^h = \mathcal{V}^h \times (\mathcal{V}^h)^{m_U} \times (\mathcal{V}^h)^{m_W} \times (\mathcal{V}^h)^{m_Q} \times (\mathcal{V}^h)^{m_R}.$$

This assumption roughly means that our splitting variables do not include flux-type variables (like $\nabla\phi$ or the normal $\frac{\nabla\phi}{|\nabla\phi|}$). The justification of this assumption can be made through appropriate inf-sup conditions on the Lagrange functionals in the above subspaces (as standard in mixed finite element discretizations, cf. [9]), at least after linearization of the system. We do not go into detail at this point for the sake of brevity, but just mention that these conditions are indeed satisfied for all the examples of flows and splitting variables we use here.

The finite element semi-discretization of the geometric flow defined via (19) is then given by the Galerkin discretization

$$\begin{aligned} \partial_\psi L[\partial_t \phi^h, W^h, Q^h; \phi^h] \eta_1 + \partial_\phi \Lambda(\phi^h, U^h, R^h) \eta_1 &= 0 & \forall \eta_1 \in \mathcal{V}^h & \quad (22) \\ \partial_W L[\partial_t \phi^h, W^h, Q^h; \phi^h] \eta_2 &= 0 & \forall \eta_2 \in (\mathcal{V}^h)^{m_U} & \quad (23) \\ \partial_Q L[\partial_t \phi^h, W^h, Q^h; \phi^h] \eta_3 &= 0 & \forall \eta_3 \in (\mathcal{V}^h)^{m_W} & \quad (24) \\ \partial_U \Lambda(\phi^h, U^h, R^h) \eta_4 &= 0 & \forall \eta_4 \in (\mathcal{V}^h)^{m_Q} & \quad (25) \\ \partial_R \Lambda(\phi^h, U^h, R^h) \eta_5 &= 0 & \forall \eta_5 \in (\mathcal{V}^h)^{m_R} & \quad (26) \end{aligned}$$

It is easy to see that the discretized problem (22)-(26) still satisfies an energy dissipation property. For this sake we choose $\eta_1 = \partial_t \phi$ and obtain

$$\partial_\psi L[\partial_t \phi^h, W^h, Q^h; \phi^h] \partial_t \phi^h = -\partial_\phi \Lambda(\phi^h, U^h, R^h) \partial_t \phi^h.$$

Using (25) und (26) we deduce

$$\begin{aligned} \frac{d}{dt} \Lambda(\phi^h, U^h, R^h) &= \partial_\phi \Lambda(\phi^h, U^h, R^h) \partial_t \phi^h + \partial_U \Lambda(\phi^h, U^h, R^h) \partial_t U^h + \\ &\quad \partial_R \Lambda(\phi^h, U^h, R^h) \partial_t R^h \\ &= \partial_\phi \Lambda(\phi^h, U^h, R^h) \partial_t \phi^h. \end{aligned}$$

Now we define the discrete energy functional

$$E_h[\varphi^h] = \Lambda(\phi^h, U^h, R^h) \tag{27}$$

with U^h and R^h determined by (25) and (26), and a discrete scalar product

$$\langle \psi_1^h, \psi_2^h \rangle_{\phi, h} = \partial_\psi L[\psi_1^h \phi^h, W^h, Q^h; \phi^h] \psi_2^h.$$

Then the semi-discrete energy dissipation becomes

$$\frac{d}{dt} E_h[\phi^h] = \frac{d}{dt} \Lambda(\phi^h, U^h, R^h) = -\langle \partial_t \phi^h, \partial_t \phi^h \rangle_{\phi, h}.$$

We finally mention that in the case of lower order energies, where we need not apply any approximation of E by a Lagrange functional, we indeed obtain $E = E_h$, i.e., energy dissipation with respect to the original energy E .

4 Time Discretization

In the following we discuss the implicit and semi-implicit time discretization of the geometric flows (19), respectively the semi-discrete problems (22)-(23). We mention again that due to the missing maximum principle these discretizations have to be interpreted in a local way, i.e., after computing one (sufficiently small) time step of the level set equation we will perform a redistancing step, with the details of redistancing to be discussed in the next section.

The time discretization will be carried out at the time steps $0 = t_0 < t_1 < \dots < t_N = T$, with local time step $\tau_k = t_{k+1} - t_k < \tau$. We shall denote the discrete solution by $\phi_k^{h, \tau} = \phi^{h, \tau}(t_k) \in \mathcal{V}^h$ (with analogous notation for the other variables).

4.1 Implicit Time Stepping

In order to gain further insight into the time discretization based on the Lagrange functionals introduced in the previous session we start with a simple implicit time discretization at least with respect to the energy functional. This means we approximate the scalar product at the previous time step, more precisely we approximate all terms depending on $\phi^{h,\tau}$ in L by values at the previous time step, and we use the backward Euler approximation

$$\partial_t \phi^{h,\tau}(t_{k+1}) \approx \frac{1}{\tau_k} (\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau}).$$

These decisions yield the implicit scheme

$$\frac{1}{\tau_k} \langle \phi_{k+1}^{h,\tau} - \phi_k^{h,\tau}, \eta_1 \rangle_k = -\partial_\phi E_h[\phi_{k+1}^{h,\tau}] \eta_1 = -\partial_\phi \Lambda[\phi_{k+1}^{h,\tau}, U_{k+1}^{h,\tau}, R_{k+1}^{h,\tau}] \eta_1$$

for all $\eta_1 \in \mathcal{V}^h$, where we use the short-hand notation

$$\langle \psi_1, \psi_2 \rangle_k := \langle \psi_1, \psi_2 \rangle_{\phi_k^{h,\tau}} = \partial_\psi L[\psi_1, W_1, Q_1; \phi_k^{h,\tau}].$$

In terms of the Lagrange functionals we can write down the fully discrete implicit scheme as the solution

$$\frac{1}{\tau_k} \partial_\phi L[\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau}, W_{k+1}^{h,\tau}, Q_{k+1}^{h,\tau}; \phi_k^{h,\tau}] \eta_1 + \partial_\phi \Lambda(\phi_{k+1}^{h,\tau}, U_{k+1}^{h,\tau}, R_{k+1}^{h,\tau}) \eta_1 = (28)$$

$$\partial_W L[\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau}, W_{k+1}^{h,\tau}, Q_{k+1}^{h,\tau}; \phi_k^{h,\tau}] \eta_2 = (29)$$

$$\partial_Q L[\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau}, W_{k+1}^{h,\tau}, Q_{k+1}^{h,\tau}; \phi_k^{h,\tau}] \eta_3 = (30)$$

$$\partial_U \Lambda(\phi_{k+1}^{h,\tau}, U_{k+1}^{h,\tau}, R_{k+1}^{h,\tau}) \eta_4 = (30)$$

$$\partial_R \Lambda(\phi_{k+1}^{h,\tau}, U_{k+1}^{h,\tau}, R_{k+1}^{h,\tau}) \eta_4 = (30)$$

for all $(\eta_1, \dots, \eta_5) \in \mathcal{P}^h$. with respect to

$$(\phi_{k+1}^{h,\tau}, W_{k+1}^{h,\tau}, U_{k+1}^{h,\tau}, Q_{k+1}^{h,\tau}, R_{k+1}^{h,\tau}) \in \mathcal{P}^h.$$

One observes that after time discretization all variations of Lagrange functionals are taken with respect to $\phi_{k+1}^{h,\tau}$ using the fixed previous time value $\phi_k^{h,\tau}$. Hence, the first equation (28) can also be reformulated as a minimization with respect to $\phi_{k+1}^{h,\tau}$. Also the other equations are optimality conditions of an optimization problem, and taking into account the usual convention for Lagrangians (minimization with respect to primal and maximization with respect to dual variables) we actually compute the variables

at the next time step $(\phi_{k+1}^{h,\tau}, W_{k+1}^{h,\tau}, U_{k+1}^{h,\tau}, Q_{k+1}^{h,\tau}, R_{k+1}^{h,\tau})$ as a solution of the saddle point problem

$$\inf_{\phi, W, U} \sup_{Q, R} \left(\frac{1}{2\tau} L[\phi - \phi_k^{h,\tau}, W, Q] + \Lambda[\phi, W, Q] \right) \quad (33)$$

over the discrete subspaces. We can also rewrite the discrete problem in the form of minimizing movements for a gradient flow (cf. [2])

$$\phi_{k+1}^{h,\tau} \in \arg \min_{\phi} \left(\frac{1}{2\tau} \|\phi - \phi_k^{h,\tau}\|_k^2 + E_h[\phi] \right),$$

with the squared norm

$$\|\phi - \phi_k^{h,\tau}\|_k^2 := \inf_W \sup_Q L[\phi - \phi_k^{h,\tau}, W, Q]$$

and

$$E_h[\phi] := \inf_U \sup_R \Lambda[\phi, U, R].$$

For illustration purpose we mention the implicit schemes for mean curvature flow

$$\int_{\Omega} \frac{(\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau})\eta}{\tau|\phi_k^{h,\tau}|} dx + \int_{\Omega} \frac{\nabla\phi_{k+1}^{h,\tau} \cdot \nabla\eta}{|\phi_{k+1}^{h,\tau}|} = 0, \quad (34)$$

and the simple surface diffusion flow

$$\begin{aligned} \int_{\Omega} Q_{k+1}^{h,\tau} \eta_1 + \int_{\Omega} \frac{\nabla\phi_{k+1}^{h,\tau} \cdot \nabla\eta}{|\phi_{k+1}^{h,\tau}|} dx &= 0 \\ \int_{\Omega} (\mathbf{P}_k \nabla(W_{k+1}^{h,\tau} + Q_{k+1}^{h,\tau})) \cdot (\mathbf{P}_k \nabla\eta_2) dx &= 0 \\ \frac{1}{\tau} \int_{\Omega} (\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau}) \eta_3 dx + \int_{\Omega} (\mathbf{P}_k \nabla Q_{k+1}^{h,\tau}) \cdot (\mathbf{P}_k \nabla\eta_3) dx &= 0, \end{aligned}$$

with $\mathbf{P}_k = \mathbf{I} - \frac{\nabla\phi_k^{h,\tau}}{|\nabla\phi_k^{h,\tau}|} \otimes \frac{\nabla\phi_k^{h,\tau}}{|\nabla\phi_k^{h,\tau}|}$. As in the continuous setting we can still deduce $Q_{k+1}^{h,\tau} = -W_{k+1}^{h,\tau}$ from the second equation and eliminate the dual variable in order to obtain an analogous form to (14), (15). For the Willmore

flow with $U = H$ we obtain

$$\begin{aligned} \int_{\Omega} \left[\frac{(\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau})\eta_1}{\tau|\phi_k^{h,\tau}|} + \frac{1}{2}U_{k+1}^{h,\tau} \frac{\phi_{k+1}^{h,\tau} \cdot \nabla\eta_1}{|\phi_{k+1}^{h,\tau}|} + \frac{\mathbf{P}_{k+1} \nabla\eta_4 \cdot \mathbf{P}_{k+1} \nabla\eta_1}{\tau|\phi_{k+1}^{h,\tau}|} \right] dx &= 0 \\ \int_{\Omega} (U_{k+1}^{h,\tau} |\phi_{k+1}^{h,\tau}| - R_{k+1}^{h,\tau}) \eta_4 dx &= 0 \\ \int_{\Omega} (-U_{k+1}^{h,\tau} \eta_5 + \frac{\nabla\phi_{k+1}^{h,\tau} \cdot \nabla\eta_5}{|\phi_{k+1}^{h,\tau}|}) dx &= 0 \end{aligned}$$

$$\text{with } \mathbf{P}_{k+1} = \mathbf{I} - \frac{\nabla\phi_{k+1}^{h,\tau}}{|\nabla\phi_{k+1}^{h,\tau}|} \otimes \frac{\nabla\phi_{k+1}^{h,\tau}}{|\nabla\phi_{k+1}^{h,\tau}|}.$$

4.2 Semi-Implicit Time Stepping

As one observes from the previous examples, the implicit schemes require the solution of strongly nonlinear systems in each time steps, which is a rather undesirable property and can be an issue with respect to computational efficiency. Indeed, the only implicit scheme in this context is the one by Chambolle (cf. [15]) for mean curvature flow, which is exactly (34) with an intermediate redistancing step. In order to overcome the difficulties with nonlinearities we consider semi-implicit time stepping, which has developed almost as a standard for numerical simulations of higher-order geometric flows, either for parametric (cf. [7, 20, 29, 30, 31, 39]), level set (cf. [24, 40]), or diffuse (cf. [8, 28]) representations of the surfaces.

In order to obtain a semi-implicit scheme with linear systems to be solved in each time step we need to construct a quadratic approximation of the energy, respectively the Lagrange functional Λ . This means we replace Λ in each iteration step by a functional $\tilde{\Lambda}^k$ quadratic with respect to ϕ and U , and linear with respect to R . The corresponding discrete energy functional is given by

$$\tilde{E}_h^k[\phi] = \inf_U \sup_R \tilde{\Lambda}^k[\phi, U, R].$$

In order to obtain a consistent approximation of the flow, the condition

$$(E_h^k)'(\phi_k^{h,\tau}) = E_h'(\phi_k^{h,\tau})$$

is needed, which can be translated into a condition on the Lagrange func-

tional as

$$\begin{aligned}\partial_\phi\Lambda(\phi_k^{h,\tau}, U_k^{h,\tau}, R_k^{h,\tau}) &= \partial_\phi\tilde{\Lambda}^k(\phi_k^{h,\tau}, \tilde{U}_k^{h,\tau}, \tilde{R}_k^{h,\tau}) \\ \partial_U\Lambda(\phi_k^{h,\tau}, U_k^{h,\tau}, R_k^{h,\tau}) &= \partial_U\tilde{\Lambda}^k(\phi_k^{h,\tau}, \tilde{U}_k^{h,\tau}, \tilde{R}_k^{h,\tau}) = 0 \\ \partial_R\Lambda(\phi_k^{h,\tau}, U_k^{h,\tau}, R_k^{h,\tau}) &= \partial_R\tilde{\Lambda}^k(\phi_k^{h,\tau}, \tilde{U}_k^{h,\tau}, \tilde{R}_k^{h,\tau}) = 0.\end{aligned}$$

Note that this condition includes the primal and dual variables ($U_k^{h,\tau}$ and $R_k^{h,\tau}$), which can be computed given $\phi_k^{h,\tau}$ through the Lagrangian Λ . These may differ from the approximating ones ($\tilde{U}_k^{h,\tau}$ and $\tilde{R}_k^{h,\tau}$), which would usually be computed from the last time step of the scheme. Hence, the derivation of a consistent scheme (and also a stable one, see below), could enforce additional computations for the computation of quadratic approximations (such as Taylor expansions). E.g., for Willmore flow one would have to compute the mean curvature variable from

$$\int_\Omega U_k^{h,\tau} \psi \, dx = - \int_\Omega \frac{\nabla \phi_k^{h,\tau} \cdot \nabla \psi}{|\nabla \phi_k^{h,\tau}|} \, dx \quad \forall \psi \in \mathcal{V}^h.$$

The next time step in a semi-implicit time scheme is then given by

$$\phi_{k+1}^{h,\tau} \in \arg \min_\phi \left(\frac{1}{2\tau} \|\phi - \phi_k^{h,\tau}\|_k^2 + \tilde{E}_h^k[\phi] \right), \quad (35)$$

i.e., as the minimizer of a quadratic optimization problem, and consequently as the solution of a linear system (the first-order optimality condition). If \tilde{E}_h^k is convex, or τ is sufficiently small, then the resulting quadratic functional is strictly convex and thus, the level set function in the next time step is unique.

For lower order energies, the construction of an approximation is again straight-forward, the approximating energy is of the form

$$\tilde{E}_h^k[\phi] = E_h[\phi_k^{h,\tau}] + E_h'[\phi_k^{h,\tau}](\phi - \phi_k^{h,\tau}) + F_h^k[\phi - \phi_k^{h,\tau}], \quad (36)$$

with a quadratic energy F_h^k . A natural choice for the quadratic term might be the second variation $F_h^k[\psi] = \frac{1}{2} E_h''[\phi_k^{h,\tau}](\psi, \psi)$, so that E_h^k would become the second-order Taylor expansion of E around the last time step. However, this choice is not essential for deriving a consistent scheme, one can instead use the appropriate quadratic functionals for increasing the stability of the scheme as we shall argue below.

For higher-order energies we construct an approximation of the Lagrangian via

$$\begin{aligned}\tilde{\Lambda}^k[\phi, U, R] &= \Lambda[\phi_k^{h,\tau}, U_k^{h,\tau}, R_k^{h,\tau}] + \\ &\quad \Lambda'[\phi_k^{h,\tau}, U_k^{h,\tau}, R_k^{h,\tau}](\phi - \phi_k^{h,\tau}, U - U_k^{h,\tau}, R - R_k^{h,\tau}) + \\ &\quad \frac{\partial^2}{\partial\phi\partial R}\Lambda[\phi_k^{h,\tau}, U_k^{h,\tau}, R_k^{h,\tau}](\phi - \phi_k^{h,\tau}, R - R_k^{h,\tau}) + \\ &\quad \frac{\partial^2}{\partial U\partial R}\Lambda[\phi_k^{h,\tau}, U_k^{h,\tau}, R_k^{h,\tau}](U - U_k^{h,\tau}, R - R_k^{h,\tau}) + \\ &\quad F_h^k[\phi - \phi_k^{h,\tau}] + G_h^k[U - U_k^{h,\tau}] - \Psi_h^k[R - R_k^{h,\tau}]\end{aligned}$$

with convex quadratic functionals F_h^k , G_h^k , and Ψ_h^k . Note that for appropriate choices of this functionals we obtain that the inf-sup (over U and R) of the approximate Lagrange functional $\tilde{\Lambda}^k$ at $\phi = \phi_k^{h,\tau}$ is attained at $\tilde{U}_k^{h,\tau} = U_k^{h,\tau} = \tilde{R}_k^{h,\tau} = R_k^{h,\tau}$, and the consistency is a direct consequence.

4.3 Stability and Energy Dissipation

Ideally we would like to obtain semi-implicit schemes that yield energy dissipation as the original geometric flows. A sufficient condition on \tilde{E}_h^k yielding energy dissipation is

$$\tilde{E}_h^k[\phi^{h,\tau}] = E_h[\phi^{h,\tau}], \quad \tilde{E}_h^k[\phi] \geq E_h[\phi] \quad \forall \phi \in \mathcal{V}^h. \quad (37)$$

Together with the definition of $\phi_{k+1}^{h,\tau}$ as a minimizer of (35) we conclude

$$\begin{aligned}\frac{1}{2\tau}\|\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau}\|_k^2 + E_h[\phi_{k+1}^{h,\tau}] &\leq \frac{1}{2\tau}\|\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau}\|_k^2 + \tilde{E}_h^k[\phi_{k+1}^{h,\tau}] \\ &\leq \tilde{E}_h^k[\phi_k^{h,\tau}] = E_h[\phi_k^{h,\tau}].\end{aligned}$$

We mention that (37) can be relaxed to

$$\frac{c}{2\tau}\|\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau}\|_k^2 + \tilde{E}_h^k(\phi) \geq E_h[\phi] \quad \forall \phi \in \mathcal{V}^h, \quad (38)$$

for some $c < 1$, which implies by an analogous argument

$$\frac{1-c}{2\tau}\|\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau}\|_k^2 + E_h[\phi_{k+1}^{h,\tau}] \leq E_h[\phi_k^{h,\tau}].$$

For a higher-order energy approximated by a Lagrangian, the first property in (37) is achieved if

$$\inf_U \sup_R \tilde{\Lambda}^k[\phi_k^{h,\tau}, U, R] = \inf_U \sup_R \Lambda[\phi_k^{h,\tau}, U, R], \quad (39)$$

which is automatic for the approximation derived in the previous section. The second (majorization) property is achieved if

$$\inf_U \sup_R \tilde{\Lambda}^k[\phi, U, R] \geq \inf_U \sup_R \Lambda[\phi, U, R] \quad \forall \phi \in \mathcal{V}^h, \quad (40)$$

whose verifications turns out to be very hard for energies like the Willmore functional.

4.4 Examples for Higher-Order Flows

In the following we provide some examples of semi-implicit schemes constructed in the above manner for some higher-order flows of particular interest:

Example 1: Mean Curvature Flow. As a first example we consider the (averaged) area functional

$$E_h[\phi] = E[\phi] = \int_{\Omega} |\nabla \phi| \, dx.$$

We look for a quadratic approximation as above with an energy of the form

$$F_h^k[\psi] = \frac{1}{2} \int_{\Omega} a_k |\nabla \psi|^2 \, dx.$$

Then the consistency and the first property of (37) are automatically satisfied. The majorization property can be rewritten as

$$\begin{aligned} \int_{\Omega} |\nabla \phi| \, dx &\leq \int_{\Omega} (|\nabla \phi_k^{h,\tau}| + \frac{\nabla \phi_k^{h,\tau} \cdot (\nabla \phi - \nabla \phi_k^{h,\tau})}{|\nabla \phi_k^{h,\tau}|} + \frac{a_k}{2} |\nabla \phi - \nabla \phi_k^{h,\tau}|^2) \, dx \\ &= \int_{\Omega} (\frac{\nabla \phi_k^{h,\tau} \cdot \nabla \phi}{|\nabla \phi_k^{h,\tau}|} + \frac{a_k}{2} |\nabla \phi - \nabla \phi_k^{h,\tau}|^2) \, dx. \end{aligned}$$

Thus, the majorization property is achieved if

$$2|\nabla \phi| |\nabla \phi_k^{h,\tau}| + 2(b_k - 1) \nabla \phi_k^{h,\tau} \cdot \nabla \phi \leq b_k |\nabla \phi|^2 + b_k |\nabla \phi_k^{h,\tau}|^2,$$

for all admissible ϕ , where we have used the notation $b_k = a_k |\nabla \phi_k^{h,\tau}|$. For $b_k \geq 1$ we obtain by the geometric-arithmetic mean inequality that

$$\begin{aligned} 2|\nabla \phi| |\nabla \phi_k^{h,\tau}| + 2(b_k - 1) \nabla \phi_k^{h,\tau} \cdot \nabla \phi &\leq 2b_k |\nabla \phi| |\nabla \phi_k^{h,\tau}| \\ &\leq b_k |\nabla \phi|^2 + b_k |\nabla \phi_k^{h,\tau}|^2, \end{aligned}$$

hence the majorization property holds. On the other hand, for $b_k < 1$ one can immediately see a violation via the choice $\phi = -\phi_k^{h,\tau}$. Hence $b_k = 1$ yields the minimal energy of the above form that satisfies the majorization property (and consequently yields stability). The resulting energy for $b_k = 1$ can be written as

$$\tilde{E}_h^k[\phi] = \frac{1}{2} \int_{\Omega} \frac{|\nabla \phi_k^{h,\tau}|^2 + |\nabla \phi|^2}{|\nabla \phi_k^{h,\tau}|},$$

and the resulting semi-implicit scheme for mean-curvature flow is

$$\int_{\Omega} \frac{(\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau})\eta}{\tau|\phi_k^{h,\tau}|} dx + \int_{\Omega} \frac{\nabla \phi_{k+1}^{h,\tau} \cdot \nabla \eta}{|\phi_k^{h,\tau}|} = 0,$$

which is exactly the one used in [18]. Also the schemes used in [19] for weighted area functionals can be derived from analogous arguments on energy approximation.

Example 2: Isotropic Surface Diffusion. As a second example we consider the surface diffusion flow with the energy equal to the area functional. We therefore use the same approximation of the energy as in the previous example

$$\tilde{E}_h^k[\phi] = \frac{1}{2} \int_{\Omega} \frac{|\nabla \phi_k^{h,\tau}|^2 + |\nabla \phi|^2}{|\nabla \phi_k^{h,\tau}|},$$

which yields unconditional stability. Together with the approximation of the H^{-1} -metric described above we arrive at the semi-implicit scheme

$$\begin{aligned} \int_{\Omega} Q_{k+1}^{h,\tau} \eta_1 dx + \int_{\Omega} \frac{\nabla \phi_{k+1}^{h,\tau} \cdot \nabla \eta}{|\phi_k^{h,\tau}|} dx &= 0 \\ \int_{\Omega} (\mathbf{P}_k \nabla (W_{k+1}^{h,\tau} + Q_{k+1}^{h,\tau})) \cdot (\mathbf{P}_k \nabla \eta_2) dx &= 0 \\ \frac{1}{\tau} \int_{\Omega} (\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau}) \eta_3 dx + \int_{\Omega} (\mathbf{P}_k \nabla Q_{k+1}^{h,\tau}) \cdot (\mathbf{P}_k \nabla \eta_3) dx &= 0, \end{aligned}$$

which is a level set analogue of the scheme used in [20] for graph representations.

Example 3: Willmore Flow As an example with an higher-order energy we consider the Willmore flow, characterized by the L^2 -metric and the energy and average energy functional, respectively,

$$e[\Gamma] = \frac{1}{2} \int_{\Gamma} h^2 dx \quad E[\phi] = \frac{1}{2} \int_{\Omega} H^2 |\nabla \phi| dx$$

with the mean curvature h and its averaged version H . Obviously we use the same approximation of the L^2 -metric as in the case of the mean curvature flow, so that it only remains to derive an energy approximation.

As noticed in Section 2, the energy functional can be calculated from the Lagrangian

$$\Lambda[\phi, U, R] = \frac{1}{2} \int_{\Omega} U^2 |\nabla \phi| \, dx + \int_{\Omega} \left(-UR + \frac{\nabla \phi \cdot \nabla R}{|\nabla \phi|} \right) \, dx.$$

We can then compute the variations

$$\begin{aligned} \partial_{\phi} \Lambda[\phi, U, R] \eta_1 &= \int_{\Omega} \left(\frac{U^2}{2} \frac{\nabla \phi \cdot \nabla \eta_1}{|\nabla \phi|} + \frac{(\mathbf{P} \nabla \eta_1) \cdot \nabla R}{|\nabla \phi|} \right) \, dx \\ \partial_U \Lambda[\phi, U, R] \eta_2 &= \int_{\Omega} (U |\nabla \phi| - R) \eta_2 \, dx \\ \partial_R \Lambda[\phi, U, R] \eta_3 &= \int_{\Omega} \left(-U \eta_3 + \frac{\nabla \phi \cdot \nabla \eta_3}{|\nabla \phi|} \right) \, dx \\ \partial^2_{\phi, R} \Lambda[\phi, U, R](\eta_1, \eta_3) &= \int_{\Omega} \frac{(\mathbf{P} \nabla \eta_1) \cdot \nabla \eta_3}{|\nabla \phi|} \, dx \\ \partial^2_{U, R} \Lambda[\phi, U, R](\eta_2, \eta_3) &= - \int_{\Omega} \eta_2 \eta_3. \end{aligned}$$

In order to complete the derivation of the semi-implicit scheme with the above quadratic form of an approximate Lagrangian $\tilde{\Lambda}^k$ it remains to specify the functionals F_h^k , G_h^k and Ψ_h^k . We shall highlight one possibility in the following, which actually yields a similar scheme as used in [24]. The idea of the choice is to look at the energy term (the first integral in Λ) for fixed U and fixed ϕ , respectively. For fixed U we obtain a weighted area functional and consequently use an analogous approximation as for area functionals above, namely

$$F_h^k[\phi] = \frac{1}{4} \int_{\Omega} (U_k^{h, \tau})^2 \frac{|\nabla \phi - \nabla \phi_k^{h, \tau}|^2}{|\nabla \phi_k^{h, \tau}|} \, dx.$$

For fixed ϕ the functional of U is already quadratic, so it seems natural to use the second-order term in the Taylor expansion

$$G_h^k[U] = \frac{1}{2} \int_{\Omega} (U - U_k^{h, \tau})^2 |\nabla \phi_k^{h, \tau}| \, dx.$$

Altogether, this yields the approximate Lagrangian

$$\begin{aligned} \tilde{\Lambda}^k[\phi, U, R] &= \frac{1}{2} \int_{\Omega} \left(U^2 |\nabla \phi_k^{h,\tau}| + (U_k^{h,\tau})^2 \frac{|\nabla \phi|^2 - |\nabla \phi_k^{h,\tau}|^2}{2|\nabla \phi_k^{h,\tau}|} \right) dx + \\ &+ \int_{\Omega} \left(-UR + \frac{(\mathbf{P}_k \nabla \phi + (\mathbf{I} - \mathbf{P}_k) \nabla \phi_k^{h,\tau}) \cdot \nabla R}{|\nabla \phi_k^{h,\tau}|} \right) dx. \end{aligned}$$

The resulting semi-implicit scheme is

$$\begin{aligned} \int_{\Omega} \left(\frac{(\phi_{k+1}^{h,\tau} - \phi_k^{h,\tau}) \eta_1}{\tau |\nabla \phi_k^{h,\tau}|} + (U_k^{h,\tau})^2 \frac{\nabla \phi_{k+1}^{h,\tau} \cdot \nabla \eta_1}{|\nabla \phi_k^{h,\tau}|} \right) dx &= 0 \\ \int_{\Omega} (U |\nabla \phi_k^{h,\tau}| - R) \eta_2 dx &= 0 \\ \int_{\Omega} \left(-U \eta_3 + \frac{(\mathbf{P}_k \nabla \phi + (\mathbf{I} - \mathbf{P}_k) \nabla \phi_k^{h,\tau}) \cdot \nabla \eta_3}{|\nabla \phi_k^{h,\tau}|} \right) dx &= 0. \end{aligned}$$

for all test functions $\eta_i \in \mathcal{V}^h$.

Example 4: Regularized Anisotropic Motion Laws. As a final example we consider the curvature regularization of an anisotropic surface energy, i.e.

$$e[\Gamma] = \int_{\Gamma} (\gamma(n) + \frac{\epsilon^2}{2} h^2) d\sigma,$$

which is of particular interest when γ (respectively its one-homogeneous extension to \mathbb{R}^d) is not convex. The averaged version of the energy is given by

$$E[\phi] = \int_{\Omega} (\gamma(\nabla \phi) + \frac{\epsilon^2}{2} H^2 |\nabla \phi|) dx.$$

In this case it is of practical interest to consider both the mean-curvature type flow (L^2 -metric) and the surface diffusion flow (H^{-1} -metric) for this energy (cf. [11, 13])

For the approximations of the L^2 -metric or H^{-1} -metric we can use exactly the same schemes as for the mean curvature or for the surface diffusion flow, respectively. Moreover, we can use the same approximation of the curvature dependent term as for the Willmore flow. Thus, we only provide an approximation of the first term in the energy

$$E_1[\phi] = \int_{\Omega} \gamma(\nabla \phi) dx.$$

A simple approximation with isotropic higher-order part is given by

$$\tilde{E}_1^k[\phi] = \int_{\Omega} \left(\gamma(\nabla\phi) + (\nabla\gamma)(\nabla\phi)(\nabla\phi - \nabla\phi_k^{h,\tau}) + \frac{\lambda}{2} \frac{(\nabla\phi - \nabla\phi_k^{h,\tau})}{|\nabla\phi_k^{h,\tau}|} \right) dx.$$

If λ is chosen large enough it is easy to see that \tilde{E}_1^k actually majorizes E_1 . A detailed discussion of semi-implicit schemes for regularized anisotropic motions can be found in [13].

5 Redistancing

It is worth noticing that for energy functionals depending on higher than first derivatives, the "global" level set approach is not well-defined. In this case the resulting evolution equation is a partial differential equation of higher than second order, and therefore does not satisfy a comparison principle. Consequently, even if a solution to the equation for the level set function ϕ exists, one cannot guarantee that ϕ is continuous and that its level sets are still the boundary of the sublevel sets (due to possible intersection and annihilation of level sets). This means that the level set approach can only be interpreted in a local sense, i.e., the level set function only satisfies the partial differential equations at the zero level set. Since one is not really interested in the other level sets of ϕ , one can use an arbitrary extension such as the signed distance function. In a theoretical approach this means one looks for a solution ϕ , which satisfies the partial differential equation only on the implicitly defined set $\{\phi = 0\}$, and is a signed distance function in the remaining part of Ω . In a practical approach one first computes one small time-steps of the partial differential equation (i.e., one violates the constraint of being a signed-distance function) and then performs a redistancing step (i.e., one computes a suitable projection to the constraint set of signed-distance functions).

As any redistancing slightly changes the interface, it should only be done if necessary. In our implementation we check the norm of the gradient of the level set function ϕ in a band around the zero level set which is about the grid size. If the norm differs too much from one, redistancing is carried out.

Our redistancing algorithm is an approach for unstructured grids based on a local Hopf-Lax formula which was introduced in [10] and given in detail for two dimensions therein. In [41] it was extended to three dimensions. In the following we give a brief outline of the idea behind and describe the algorithm in detail. Let $\tilde{\phi}$ be a level set function which gives implicitly

the interface Γ , i.e. $\Gamma = \{x \in \Omega \mid \tilde{\phi}(x) = 0\}$, and $\Omega^+ = \{x \in \Omega \mid \tilde{\phi}(x) > 0\}$, $\Omega^- = \{x \in \Omega \mid \tilde{\phi}(x) < 0\}$ denote the domain with positive and negative level sets respectively. We want to construct a signed distance function with respect to the zero level set of $\tilde{\phi}$. It is sufficient to calculate a distance function. The sign then can easily be added with the help of $\tilde{\phi}$.

The eikonal equation

$$\begin{aligned} |\nabla u| &= 1 \quad \text{on } \Omega \\ u|_{\partial\Omega} &= g \end{aligned}$$

with adequate function g is a Hamilton-Jacobi equation of the type

$$H(x, \nabla u) = 1 \quad \text{on } \Omega \tag{41}$$

$$u|_{\partial\Omega} = g. \tag{42}$$

It can be shown that under certain assumptions on H and g a solution of (41), (42) is given by the Hopf-Lax formula. For the eikonal equation where $H(\nabla u) = |\nabla u| - 1$ the Hopf-Lax formula reads

$$u(x) = \min_{y \in \partial\Omega} (g(y) + |x - y|).$$

The principle idea of the redistancing algorithm is to apply the Hopf-Lax formula on a local discretized problem.

Let Ω_h be a polygonal approximation of Ω , \mathcal{T}_h the corresponding triangulation and \mathcal{V}^h the set of linear finite element functions on Ω_h . We denote \mathcal{T}_h^I the set of elements $\sigma \in \mathcal{T}_h$ which are cut by the interface ("interface elements"), i.e.

$$\mathcal{T}_h^I = \{\sigma \in \mathcal{T}_h \mid \sigma \cap \Gamma_h \neq \emptyset\},$$

with Γ_h the discretized interface given implicitly by the level set function $\phi_h \in \mathcal{V}^h$, and

$$\Omega_h^I = \bigcup_{\sigma \in \mathcal{T}_h^I} \sigma.$$

We are looking for the solution of the discretized problem

$$|\nabla \phi_h| = 1 \quad \text{on } \Omega_h \tag{43}$$

$$\phi_h|_{\partial\Omega_h^I} = g_h, \tag{44}$$

where for a vertex x_h the value $g_h(x_h)$ is the distance of x_h to the discretized interface Γ_h . For each vertex $x_h \notin \Omega_h^I$ and its local element patch $\omega_h(x_h)$

which is defined as the interior of the union of all elements $\sigma \in \mathcal{T}_h \setminus \mathcal{T}_h^I$ containing x_h , a solution of the local discretized problem

$$\begin{aligned} |\nabla \phi_h^*| &= 1 \quad \text{on } \omega_h(x_h) \\ \phi_h^*|_{\partial\omega_h(x_h)} &= g_h \end{aligned}$$

is given by the local Hopf-Lax formula

$$\phi_h^*(x_h) = \min_{y \in \partial\omega_h(x_h)} (g_h(y) + |x_h - y|).$$

The solution of (43), (44) results now from an iterative application of the local Hopf-Lax formula for the vertices x_h of the triangulation. We define the local Hopf-Lax update by

$$(\Lambda_h \phi_h)(x_h) = \begin{cases} \min_{y \in \partial\omega_h(x_h)} (\phi_h(y) + |x_h - y|) & \text{for } x_h \notin \Omega_h^I \\ \phi_h(x_h) & \text{for } x_h \in \Omega_h^I. \end{cases}$$

A possible redistancing algorithm consists in traversing all vertices x_h until the Hopf-Lax updates do not give new values or to be more precise until for any small tolerance tol we have $|(\Lambda_h \phi_h)(x_h) - \phi_h(x_h)| \leq tol$ for all vertices x_h . Because our approach is a slight variation of this idea, we give a short explanation for the convergence of the Hopf-Lax updates which is based on a monotonicity argument: The sequence $(u_h^n)_{n \geq 0}$ defined through

$$\phi_h^0(x_h) = \begin{cases} \infty & \text{for } x_h \notin \Omega_h^I \\ g_h(x_h) & \text{for } x_h \in \Omega_h^I \end{cases}$$

and

$$\phi_h^{n+1} = \Lambda_h \phi_h^n$$

is decreasing monotonously,

$$\phi_h^{n+1} \leq \phi_h^n \leq \dots \leq \phi_h^1 \leq \phi_h^0.$$

As for each vertex x_h there is the lower bound

$$\phi_h^n(x_h) \geq \min_{y_h \in \Omega_h^I} g_h(y_h) \quad \text{for all } n \geq 0,$$

the sequence converges uniformly to a function ϕ_h .

Our approach uses an element based Hopf-Lax vertex update

$$(\Lambda_h \phi_h)(x_h, \sigma) = \begin{cases} \min_{y \in \partial(\omega_h(x_h) \cap \sigma)} (\phi_h(y) + |x_h - y|) & \text{for } x_h \notin \Omega_h^I \\ \phi_h(x_h) & \text{for } x_h \in \Omega_h^I. \end{cases} \quad (45)$$

We make use of the following redistancing algorithm:




Figure 1: 3d-Hopf-Lax vertex update algorithm: Shows the rotation of an element with respect to ϕ_h (on the left) and restrictions to the existence of this rotation (on the right).

1. For all vertices x_h in Ω_h^I :
 Calculate the distance $g_h(x_h)$ to the discretized interface.
- 2.

$$\phi_h(x_h) = \begin{cases} \infty & \text{for } x_h \notin \Omega_h^I \\ g_h(x_h) & \text{for } x_h \in \Omega_h^I \end{cases}$$

3. Do
 For all grid elements σ :
 For all vertices x_h in σ :
 If $(|(\Lambda_h \phi_h)(x_h) - \phi_h(x_h)| > tol)$
 $\phi_h(x_h) = (\Lambda_h \phi_h)(x_h)$.
 While (Hopf-Lax vertex update has been performed during grid
 traverse).

The algorithm finishes after a finite number of iterations because the monotonicity property is maintained. In the rest of the chapter we describe in detail the algorithm for the calculation of the Hopf-Lax vertex update (45). It only makes use of geometric considerations. For dimensions two, it is quite short, the implementation only takes a few lines. The algorithm is given in [10]. We now describe an algorithm for dimensions three.

We denote $HL\text{-}2d(x_h, \sigma)$, $HL\text{-}3d(x_h, \sigma)$ the Hopf-Lax vertex update for the vertex x_h in element σ for dimensions two and dimensions three respectively. To explain the algorithm we employ a geometric interpretation of the Hopf-Lax vertex update $HL\text{-}3d(x_h, \sigma)$. Denote u_h, v_h, w_h the vertices in σ in the face opposite to x_h . Now we place σ in three-dimensional space such




Figure 2: 3d-Hopf-Lax vertex update algorithm: Possible positions of the projection point D with respect to \triangle_{ABC} .

$$\begin{aligned}
D \in P_{ABC} : & \quad HL_3d(x_h, \sigma) = \text{distance of } x_h \text{ to x-y-plane} \\
D \in P_A : & \quad HL_3d(x_h, \sigma) = \phi_h(u_h) + |x_h - u_h| \\
D \in P_B : & \quad HL_3d(x_h, \sigma) = \phi_h(v_h) + |x_h - v_h| \\
D \in P_C : & \quad HL_3d(x_h, \sigma) = \phi_h(w_h) + |x_h - w_h| \\
D \in P_{AB} : & \quad HL_3d(x_h, \sigma) = HL_2d(x_h, \triangle_{x_h u_h v_h}) \\
D \in P_{AC} : & \quad HL_3d(x_h, \sigma) = HL_2d(x_h, \triangle_{x_h u_h w_h}) \\
D \in P_{BC} : & \quad HL_3d(x_h, \sigma) = HL_2d(x_h, \triangle_{x_h v_h w_h})
\end{aligned}$$

Table 1: 3d-Hopf-Lax vertex update algorithm: Calculation formula for the Hopf-Lax update depending on the position of D (see Figure 2).

that u_h has the height $\phi_h(u_h)$ over the x-y-plane, v_h the height $\phi_h(v_h)$ and w_h the height $\phi_h(w_h)$. That means the x-y-plane serves as an approximation of the interface, see the left picture in figure 1. If the perpendicular line from the rotated x_h to the x-y-plane passes through $\triangle_{u_h v_h w_h}$ the Hopf-Lax vertex update is the distance of x_h to the x-y-plane. In other cases we need further geometric arguments. Denote A the projection of u_h on the x-y-plane, B the projection of v_h , C the projection of w_h and D the projection of x_h . Then the position of D can be described by the areas defined in figure 2. Let $y(x_h)$ be defined through

$$y(x_h) = \operatorname{argmin}_{y \in \partial(\omega_h(x_h) \cap \sigma)} (g_h(y) + |x_h - y|).$$

The position of D gives the position of $y(x_h)$. For example, if D is situated in the area P_{AB} , the point $y(x_h)$ lies on the edge of σ containing u_h and v_h . That means the Hopf-Lax vertex update is obtained as the two-dimensional Hopf-Lax vertex update on a face of σ , namely

$$HL_3d(x_h, \sigma) = HL_2d(x_h, \triangle_{x_h u_h v_h}).$$

Or, if D lies in the area P_C , we have $y(x_h) = w_h$. Thus depending on the position of D , the three-dimensional Hopf-Lax vertex update can be calculated as in table 1. Our algorithm now consists in first performing the above rotation with respect to the distances $\phi_h(u_h)$, $\phi_h(v_h)$ and $\phi_h(w_h)$, and then calculating the update as in table 1.

In some cases the rotation is not possible. Let us consider the rotation to be done in three steps: First we arrange the vertices u_h , v_h and w_h such that

$$\phi_h(u_h) \leq \phi_h(v_h) \leq \phi_h(w_h). \quad (46)$$

Then the edge $[u_h, v_h]$ is rotated appropriately. This is only possible if

$$\phi_h(v_h) - \phi_h(u_h) \leq |v_h - u_h|. \quad (47)$$

And finally we position the edge $[u_h, w_h]$. The right picture in figure 1 shows a rotation of $\Delta_{u_h v_h w_h}$ into the x-z-plane and illustrates that the second rotation is only possible if

$$0 \leq \alpha + \beta \leq \pi/2 \quad (48)$$

$$\frac{\phi_h(w_h) - \phi_h(u_h)}{|w_h - u_h|} \leq \sin(\alpha + \beta), \quad (49)$$

with the angles α and β defined in the figure. The algorithm for the three-dimensional Hopf-Lax vertex update $HL_3d(x_h, \sigma)$ is now as follows:

1. Arrange u_h , v_h and w_h as in (46).
2. If (rotation of σ is possible, i.e. (47)-(49) are fulfilled)
 Rotate σ and calculate $HL_3d(x_h, \sigma)$ as in table 1.
 Else
 $HL_3d(x_h, \sigma)$ is the minimum of the two-dimensional Hopf-Lax vertex updates for x_h on the faces of σ containing x_h .

The described algorithm is well suited for unstructured grids. Further speed up can be achieved by an ordered element travers, in which elements near the interface are visited first.

6 Implementation and results

The derived numerical schemes are implemented in the adaptive finite element toolbox AMDiS [4]. The toolbox provides a framework for the efficient solution of systems of partial differential equations by adaptive finite

elements on unstructured grids. For details on the software we refer to [42, 41, 43].

In order to reduce the computational overhead associated with the level set method we use adaptive mesh refinement and coarsening. Thus only within a small neighborhood (narrow band) around $\Gamma(t) = \{\phi(\cdot, t) = 0\}$ we allow for a fine resolution of the grid. Whereas away from the zero level set the grid is coarsened. As an indicator for refinement and coarsening serves the signed distance function.

In all simulations we use piecewise linear finite elements and solve the resulting linear system using a Schur complement and a GMRES solver.

As examples we consider variants of the H^{-1} -gradient flow of the energy

$$e_1[\Gamma] = \int_{\Gamma} \gamma \, d\sigma, \quad \text{and} \quad e_2[\Gamma] = \int_{\Gamma} h^2 \, d\sigma,$$

with $\gamma = \gamma(n)$. Fig. 3 shows the evolution of a cube to a sphere by surface diffusion ($\gamma = 1$). It is an example for a fourth order equation. The discretization is described in detail in Section 4.

As an example of a sixth order equation we use a combination of the energies e_1 and e_2

$$e_3[\Gamma] = \int_{\Gamma} \gamma + \frac{\epsilon^2}{2} h^2 \, d\sigma,$$

with ϵ a small parameter. Such energies arise in materials science with γ being non-convex [23, 33, 27, 38]. Thus the curvature term can be viewed as a regularization. For an approach to discretize the equations resulting from the anisotropic term we refer to [13]. The discretization of the remaining terms is described in detail in Section 4. Fig. 4 shows the evolution of a cube to the Wulff shape by curvature regularized anisotropic surface diffusion.

Acknowledgments

A.V. and C.S. have been supported by the German Science Foundation DFG through project SFB 611 / B3. M.B. and A.V. acknowledge partial support by the NSF-funded Institute for Pure and Applied Mathematics (IPAM), UCLA, part of this work has been performed during their participation in the IPAM program *Bridging Time and Length Scales in Materials Science and Bio-Physics*.




Figure 3: Evolution by surface diffusion (isotropic evolution). Simulation parameters: $[0, 4] \times [0, 4] \times [0, 4]$ grid, grid size $h = 0.05$, timestep $\Delta t = 10^{-6}$. From top left to bottom right: $t = 0.0, 0.001, 0.005, 0.01$.




Figure 4: Evolution by curvature regularized surface diffusion (anisotropic evolution). Simulation parameters: $[0, 4] \times [0, 4] \times [0, 4]$ grid, grid size $h = 0.05$, timestep $\Delta t = 10^{-6}$. From top left to bottom right: $t = 0.0, 0.001, 0.005, 0.01$.

References

- [1] F.Almgren, J.E.Taylor, *Optimal geometry in equilibrium and growth*, *Fractals* **3** (1996), 713-723.
- [2] L.Ambrosio, N.Gigli, G.Savare, *Metric Gradient Flows* (Birkhäuser, Basel, 2005).
- [3] L.Ambrosio, N.Fusco, D.Pallara, *Functions of bounded variation and free discontinuity problems.*, Oxford University Press (2000)
- [4] www.caesar.de/amdis.html
- [5] M.Bertalmio, L.T.Cheng, S.Osher, G.Sapiro, *Variational problems and partial differential equations on implicit surfaces: The framework and examples in image processing and pattern formation*, *J. Comput. Phys.* **174** (2001), 759-780.
- [6] M.Bertalmio, F.Memoli, L.T.Cheng, G.Sapiro, S.Osher, *Variational problems and partial differential equations on implicit surfaces: Bye bye triangulated surfaces ?*, in: S.Osher, N.Paragios, eds., *Geometric Level Set Methods in Imaging, Vision, and Graphics* (Springer, New York, 2003), 381-398.
- [7] E.Bänsch, P.Morin, R.H.Nochetto *A finite element method for surface diffusion: the parametric case*, *J. Comput. Phys.* **203** (2005), 321-343
- [8] J.W. Barrett, J.F. Blowey, *Finite element approximation of the Cahn-Hilliard equation with concentration dependent mobility*, *Math. Comp.* **68** (1999) 487517.
- [9] F.Brezzi, M.Fortin, *Mixed and Hybrid Finite Element Methods*, (Springer, New York, 1991).
- [10] F.Bornemann, C.Rasch, *Finite-Element discretization of static Hamilton-Jacobi equations based on a local variational principle* *Comput. Vis. Sci.* **9** (2006), 57-69.
- [11] M.Burger, *Numerical simulation of anisotropic surface diffusion with curvature-dependent energy*, *J. Comp. Phys.* **203** (2005), 602-625.
- [12] M.Burger, *Finite element approximation of elliptic partial differential equations on implicit surfaces*, CAM-Report 05-46 (UCLA, 2005).

- [13] M.Burger, F.Hausser, C.St"ocker, A.Voigt, *A level set approach to anisotropic flows with curvature regularization*, J. Comp. Phys. (2007), to appear.
- [14] W.C.Carter, J.W.Cahn, J.E.Taylor, *Variational methods for microstructural evolution*, JOM **49** no. 12 (1998), 30-36.
- [15] A.Chambolle, *An algorithm for mean curvature motion* Interfaces Free Bound. **6** (2004), 195-218.
- [16] D.Chopp, J.A.Sethian, *Motion by intrinsic Laplacian of curvature*, Interfaces and Free Boundaries **1** (1999), 107-123.
- [17] U.Clarenz, F.Haußer, M.Rumpf, A.Voigt, U.Weikard, *On level set formulations for anisotropic fourth order geometric evolution problems*, in A.Voigt, ed., *Multiscale Modeling in Epitaxial Growth* ISNM 149 (Birkhäuser, Basel, 2005), 227-237.
- [18] K.Deckelnick, G.Dziuk, *Error estimates for a semi implicit fully discrete finite element scheme for mean curvature flow of graphs*, Interfaces Free Boundaries **2** (2000), 341-359.
- [19] K.Deckelnick, G.Dziuk, *A fully discrete numerical scheme for weighted mean curvature flow*, Numer. Math. **91** (2002), 423-452.
- [20] K.Deckelnick, G.Dziuk, C.M.Elliott, *Fully discrete semi-implicit second order splitting for anisotropic surface diffusion of graphs*, Preprint (Isaac Newton Institute, Cambridge, 2003).
- [21] K.Deckelnick, G.Dziuk, C.M.Elliott, *Computation of geometric partial differential equations and mean curvature flow*, Acta Numerica (2005), 139-232.
- [22] M.C.Delfour, J.P.Zolésio, *Shapes and geometries. Analysis, differential calculus, and optimization* (SIAM, Philadelphia, 2001).
- [23] A.DiCarlo, M.Gurtin, P.Podio-Guidugli, *A regularized equation for anisotropic motion by curvature*, SIAM J. Appl. Math. **52** (1992), 1111-1119.
- [24] M.Droske, M.Rumpf, *A level set formulation for Willmore flow*, Interfaces and Free Boundaries **6** (2004), 361-378.
- [25] Q.Du, C.Liu, R.Ryham, X.Wang, *A phase field formulation of the Willmore problem.*, Nonlinearity **18** (2005), 1249-1267.

- [26] L.C.Evans, R.F.Gariepy, *Measure Theory and Fine Properties of Functions*, CRC Press (1992)
- [27] E.Fried, M.E.Gurtin, *A unified treatment of evolving interfaces accounting for small deformations and atomic transport with emphasis on grain-boundaries and epitaxy*, Adv. Appl. Mech. **40** (2004), 1-177.
- [28] K.Glasner, *A diffuse interface approach to HeleShaw flow*, Nonlinearity **16** (2003), 4966.
- [29] F.Haußer, A.Voigt, *A discrete scheme for regularized anisotropic surface diffusion, a sixth order geometric evolution equation*, Interf. Free Bound. **7** (2005), 1-17
- [30] F.Haußer, A.Voigt, *A discrete scheme for regularized anisotropic curve shortening flow*, Appl. Math. Lett. **19** (2006), 691-698
- [31] F.Haußer, A.Voigt, *A Discrete Scheme for Parametric Anisotropic Surface Diffusion*, J. Sci. Comput. (2006), in press
- [32] C.Herring, *Some theorems on the free energies of crystal surfaces*, Phys. Rev. **82** (1951), 87-93.
- [33] M.E.Gurtin, M.E.Jabbour, *Interface evolution in three dimensions with curvature-dependent energy and surface diffusion: Interface-controlled evolution, phase transitions, epitaxial growth of elastic films*, Arch. Rat. Mech. Anal. **163** (2002), 171-208
- [34] W.W.Mullins, *Two-dimensional motion of idealized grain boundaries*, J. Appl. Phys. **27** (1956), 900-904.
- [35] W.W.Mullins, *Theory of thermal grooving*, J. Appl. Phys. **28** (1957), 333-339.
- [36] S.J.Osher, R.P.Fedkiw, *The Level Set Method and Dynamic Implicit Surfaces* (Springer, New York, 2002).
- [37] S.J.Osher, J.A.Sethian, *Fronts propagating with curvature-dependent speed: Algorithms based on Hamilton–Jacobi formulations*, J. Comp. Phys., **79** (1988), 12-49.
- [38] A.Rätz, A.Voigt, *Higher order regularization of anisotropic geometric evolution equations in three dimensions* J. Comput. Theor. Nanosci. **3** (2006), 543-560.

- [39] R.E. Rusu, *An algorithm for the elastic flow of surfaces*, Interfaces and Free Boundaries **7** (2005), 229-239.
- [40] P.Smereka, *Semi-implicit level set methods for curvature and surface diffusion motion*, J. Sci. Comp. **19** (2003), 439-456.
- [41] C.Stöcker, S.Vey, A.Voigt, *AMDiS - Adaptive multidimensional simulations: composite finite elements and signed distance functions*, WSEAS Trans. Circ. Syst. **4** (2005) 111-116.
- [42] S.Vey, A.Voigt, *AMDiS - Adaptive Multidimensional Simulations: Object oriented software concepts for scientific computing* WSEAS Trans. Syst **3** (2004), 1564-159
- [43] S.Vey, A.Voigt, *AMDiS - Adaptive Multidimensional Simulations* Comp. Vis. Sci., in press