NUMERICAL METHODS FOR ANISOTROPIC MEAN CURVATURE FLOW BASED ON A DISCRETE TIME VARIATIONAL FORMULATION. *

ADAM OBERMAN [†], STANLEY OSHER [‡], RYO TAKEI [§], AND RICHARD TSAI [¶]

Abstract. Numerical methods for planar anisotropic mean curvature flow are presented for smooth and crystalline anisotropies. Early numerical work by J. Taylor was based on directly minimizing the surface free energy using parameterized curves. The level set method approach of Osher and Sethian can be used in the case of smooth anisotropies but not in the crystalline case, where the motion is nonlocal. A. Chambolle combined the level set approach in the variational framework of Almgren, Taylor and Wang, which results in a total variation minimization problem. Our approach uses the split Bregman method for the total variation minimization. This method has the advantage of decoupling the anisotropy case, we derive an explicit formula for the crystalline shrinkage operator. In the smooth anisotropy case, the explicit shrinkage operator is not available: instead we solve a system of nonlinear equations, called inverse scale space flow, at each iteration. Numerical results are presented.

Key words.

subject classifications.

1. Introduction We study numerical methods for the time evolution of a closed planar curve C(t) with normal velocity equal to its anisotropic mean curvature. The motion is the gradient flow of the anisotropic perimeter

$$\mathcal{E}_{\phi}(C) = \int_{C} \phi(\hat{n}) \, ds. \tag{1.1}$$

Here, \hat{n} is the unit outer normal to the curve *C* and *ds* is the arc-length measure. For a derivation, we refer to the survey [Tay96] and the references therein. In the case $\phi(\cdot) = |\cdot|$, the motion is a natural generalization to the well known (isotropic) mean curvature flow, or the curve-shortening flow, where the curve evolves to minimize its length.

The direct approach is to solve the gradient flow of the surface free energy for a given parameterized curve. The formulation is as follows: let $\eta(C)$ be a small deformation of a closed curve C. Then, for a 'time step' h > 0, find the particular deformation η^* that minimizes

$$\min_{\eta} \mathcal{E}_{\phi}(\eta(C)) - \mathcal{E}_{\phi}(C) + \frac{1}{2h} \langle \eta, \eta \rangle.$$
(1.2)

It can be shown (c.f. [Tay96]) that, $\eta^*(C)$ is an approximation to the anisotropic mean curvature evolution of C by time h. This variational approach was studied in [ATW93, LS95, CN07].

The level set method (see Section 1.2) can also be used to implement isotropic and smoothly anisotropic motions. It cannot be used in the crystalline case because the motion is nonlocal. However, it is possible to combine the advantages of the

1

^{*}Received date / Revised version date

[†]Department of Mathematics, Simon Fraser University, (aoberman@sfu.ca).

[‡]Department of Mathematics, University of California, Los Angeles, (sjo@math.ucla.edu).

[§]Department of Mathematics, University of California, Los Angeles, (rrtakei@ucla.edu).

[¶]Department of Mathematics, University of Texas, Austin, (ytsai@math.utexas.edu).

Anisotropic Mean Curvature Flow

level set approach with the variational formulation. Chambolle [Cha04a] showed that minimizing (1.2) can be achieved by defining f to be the signed distance function to the curve C, and performing the minimization

$$\min_{u \in L^{2}(\Omega)} \int_{\Omega} \phi(\nabla u) + \frac{1}{2h} ||u - f||^{2}_{L^{2}(\Omega)}$$
(1.3)

The first term is the anisotropic total variation, see the definition (2.5) below. If u^* is the minimizer of (1.3), then $\eta^*(C) = \{u^*(x) = 0\}$, the zero level set of u^* .

1.1. Contribution of this work In this article, we consider two different scenarios of (1.3): one involving a non-smooth ϕ and the other with smooth but anisotropic ϕ . Consequently, we propose and implement two numerical algorithms for these two scenarios. The first numerical method treats crystalline anisotropies, and takes advantage of an explicit formula we derive in this case. Crystalline anisotropies are of interest, for example, in crystal growth, c.f. [Gur93]. The second numerical method treats the case of smooth anisotropies and yields a theoretically interesting result relating the level set and variational approaches.

To alleviate the difficulty in solving the nonlinear (and singular in the case of crystalline curvatures) minimization that arises from (1.3), we propose to deal with it by extensions of the split Bregman method, which solves (1.3) via a sequence of easier-to-solve problems. The crystalline anisotropy case is treated by modifying the split Bregman based algorithm [GO09] applied to isotropic mean curvature flow: in particular, we characterize the solution of the anisotropic shrinkage (or soft thresholding) problem: given $y \in \mathbb{R}^2$, find

$$\arg\min_{x \in \mathbb{R}^2} \phi(x) + \frac{1}{2} |x - y|^2.$$
 (1.4)

In Theorem 4.3, we derive a formula for the minimizer of (1.4); embedding this formula within the split Bregman algorithm gives rise to the corresponding anisotropic mean curvature flow. The main advantage of this approach is that it circumvents the difficulty of treating the non-smoothness of ϕ by confining the anisotropy only within the subproblem (1.4); this observation was mentioned briefly in [GBO09], but an explicit characterization is novel. While our solution to (1.4) is simple, an analytically explicit formula for general (e.g. smooth) ϕ is unavailable. Due to the convexity of the problem, one may employ efficient numerical methods to solve (1.4); we will comment on this later in Section 5.

For the smooth anisotropy case, we propose to solve a system of nonlinear equations, called an *inverse scale space flow*. This is inspired by the work of Burger et al. [BGOX06], where they interpret the iterative refinement of the Rudin-Osher-Fatemi (ROF) image denoising scheme as a discretization of a continuous evolution equation. In their context, the evolution initially captures large scale artifacts of a given noisy image f, then gradually finer scales are captured, eventually converging to f. For the present mean curvature problem, we apply the same strategy but to the split Bregman functional (5.5); to our knowledge, this application is new. The resulting inverse scale space flow share several properties to that of Burger et al. (e.g. decay of Bregman distance, result 6 of Lemma 5.3), but is distinct in that it *evolvess* two functions that converge to each other in the limit. It is also of theoretical interest that the (nonlinear) evolution equations involves the linear term Δu , rather than the degenerate elliptic curvature div $(\nabla u/|\nabla u|)$, a favorable property inherited from the split Bregman framework. We also show that the limit of the evolution is in fact the update solution to a semi-implicit Euler scheme for the mean curvature PDE, see remark 5.1. Convergence proofs and numerical results are given.

For a brief outline of the proposed method in the context of the split Bregman method, see Section 2.2.

1.2. Motion of level sets by mean curvature An alternative method to the direct minimization of the surface free energy described above is to use the level set approach due to Osher and Sethian [OS88]. The advantages of the level set approach are well documented: it handles topology changes and can be implemented on a uniform grid. Note, however, for non-smooth surface free energies, such as the crystalline case, the level set method cannot be used, since the motion is nonlocal [ATW93]. (See (1.9), the term $\gamma(\omega) + \gamma''(\omega)$ is undefined in the crystalline case).

Represent the curve $C(t) \subset \mathbb{R}^2$ as the zero level set of an auxiliary function u(x,t), $x \in \mathbb{R}^2$. Then, the idea is to evolve u by a degenerate elliptic partial differential equation (PDE)

$$\frac{\partial u}{\partial t} = |\nabla u| \operatorname{div} \frac{\nabla u}{|\nabla u|}.$$
(1.5)

In fact, all level curves $\{x \in \mathbb{R}^2 : u(x,t) = \text{constant}\}\$ evolve by mean curvature under (1.5). This PDE was analyzed in the viscosity solutions framework in [ES91, CGG91].

The anisotropic analogue of (1.5) may take various forms. The general form is,

$$\frac{\partial u}{\partial t} = \beta \left(\frac{\nabla u}{|\nabla u|} \right) |\nabla u| \operatorname{div} \nabla \phi(\nabla u).$$
(1.6)

where β is called the *mobility function*. It has been shown [GL94, DGM96] that this PDE produces self-similar solutions in two dimensions for positive, continuous β . Two special cases have been studied previously: the variational approach of Almgren, Taylor and Wang [ATW93] solved the case $\beta = 1$, while Chambolle [Cha04a] and Novaga and Paulini [NP07] studied the case $\beta = \gamma$, where $\gamma(x) = \phi(x)/|x|$. We chose to describe our methods for the anisotropic form

$$\frac{\partial u}{\partial t} = |\nabla u| \operatorname{div} \nabla \phi(\nabla u), \tag{1.7}$$

i.e. the case $\beta = 1$, mainly to simplify the implementation, but other choices for the mobility function can be implemented by the present algorithms similarly, see remarks 4.2 and 5.2.

Let us write,

$$\phi(\nabla u) = \gamma(\omega) |\nabla u|, \qquad \omega = \tan^{-1}(u_y/u_x). \tag{1.8}$$

If γ is smooth, (1.7) can be written as

$$\frac{\partial u}{\partial t} = (\gamma(\omega) + \gamma''(\omega)) |\nabla u| \operatorname{div} \frac{\nabla u}{|\nabla u|}.$$
(1.9)

The condition $\gamma(\omega) + \gamma''(\omega) > 0$ to avoid ill-posedness is a standard assumption in interface motion, c.f. [Gur93, Chapter 9]. The PDE (1.9) for smooth ϕ is still solvable numerically, for example, using conventional, explicit finite difference schemes. We will see in Section 5.3 (Proposition 5.2) that the inverse scale space flow approximates a discretization of (1.9) in t.

Anisotropic Mean Curvature Flow

1.3. Related work In this Section we discuss related work.

Theoretical work on anisotropic mean curvature flow have been studied extensively; we refer the interested reader to the survey article of Bellittini [Bel04] (on crystalline mean curvature flows), the book by Giga [Gig06] (on various interface problems, in the context of the level set equations) and the references therein.

Several numerical methods for Chambolle's approach have been studied. In the case $\phi = |\cdot|$, (1.3) is the Rudin-Osher-Fatemi (ROF) functional used for image denoising [ROF92]. Numerical methods based in the variational approach to mean curvature flow has recently gained attention due to the advent of fast ROF (minimization) solvers combined with efficient methods for computing the distance function. Chambolle and Darbon [CD09] used a combinatorical ROF solver, referred to as 'graph-cuts', to solve anisotropic mean curvature flow; the algorithm is designed only for crystalline ϕ with the singularities aligned along grid directions. Chambolle [Cha04a] also proposed a ROF solver [Cha04b], a fixed point method of a projection in the dual space, and solved both isotropic and crystalline anisotropic cases (in the latter case, numerical results were presented without proof of convergence). For the crystalline case, the present paper follows [GB009] where the split Bregman method [G009] is used to solve the ROF minimization.

Numerical methods for the direct solution of the level set PDE for mean curvature flow have also been studied. Monotone, convergent finite difference schemes for the isotropic mean curvature motion PDE (1.5) were discovered by Catte, et al. [CDK95] and Oberman [Obe04]. Clarenz, et al. [CHR⁺05] considered a weak formulation of a (ϵ -regularized) level set equation for anisotropic mean curvature motion and solved it using a finite element scheme; later this technique was generalized to higher orer flows in [BSV08]. See also the survey article [DDE05] for a semi-implicit Euler scheme of the level set PDE, implemented using the finite element method.

The Merriman-Bence-Osher scheme [MBO94] is a theoretical method for approximating isotropic motion by mean curvature. It approximates a mean curvature flow by repeatedly solving the heat equation for a short time and thresholding. Generalization to anisotropic flow have been studied in [CN06]. In a series of papers, [ET06, ERT05, ERT08, ERT10], Esedoglu et al. proposed a new class of algorithms for the computation of geometric motions of interfaces, including isotropic curvature motions and higher order motions such as motion by surface diffusion. These algorithms can be seen as variants of Merriman, Bence, and Osher's threshold dynamics, and consist of two alternating steps: the construction of either a characteristic function or a signed distance function to the interface, and convolution with a suitable kernel. In particular, by using signed distance functions, the algorithms proposed in [ERT10] allow for high accuracy on uniform grids. Furthermore, motion by smooth anisotropic curvatures can also be computed after a slight modification of the first step. These algorithms preserve the computational efficiency of the Merriman-Bence-Osher scheme; in comparison to the conventional level set methods, they avoid the evaluation of degenerate nonlinear PDEs; in comparison to Chambolle's algorithm, they avoid the computation of nontrivial optimization problem. In this regard, the present article's contribution lies in alleviating the complexity for solving the nontrivial optimization problems in the framework on Chambolle. There has been other generalizations of the MBO scheme, involving the use of characteristic functions, see e.g. [Ruu98, RM00], and [GH08] for flows involving fourth-order partial derivatives.

Hausser and Voigt [HV06] considered a regularized mean curvature flow by adding $\epsilon \kappa^2$ (κ is curvature and $\epsilon \ll 1$) to the interfacial energy (1.1); the evolution of the

interface in two dimensions was solved via a marker-point type scheme, c.f. [Set99, II.3.1].

2. Definitions and outline of main algorithms

2.1. Definitions and notations Throughout this article, Ω will be a bounded, connected, open subset of \mathbb{R}^2 .

We will start by defining ϕ , the surface free energy function, (using a term from thermomechanics [Gur93]). Assume

 ϕ is a convex, even, positively 1-homogeneous function with $0 < a(x) \le \phi(x) \le A(x) < \infty$, for all $x \in \mathbb{R}^2$.

The unit ball of the surface free energy function is known as the *Frank diagram*:

$$\mathcal{F}_{\phi} := \{ y \in \mathbb{R}^2 \, | \, \phi(y) \le 1 \}.$$
(2.1)

Also define ϕ° , the dual norm (or polar) of ϕ :

$$\phi^{\circ}(x) := \sup_{\phi(y) \le 1} x \cdot y. \tag{2.2}$$

The Wulff shape of ϕ is the unit ball of ϕ° :

(

$$\mathcal{W}_{\phi} := \left\{ y \in \mathbb{R}^2 : \phi^{\circ}(y) \le 1 \right\}, \tag{2.3}$$

Note that, by the convexity of ϕ , the Wulff shape \mathcal{W}_{ϕ} is convex. In addition, the Wulff shape and Frank diagram are related by duality:

$$\phi(x) = \sup_{\phi^{\circ}(y) \le 1} y \cdot x = \sup_{y \in \mathcal{W}_{\phi}} y \cdot x.$$
(2.4)

EXAMPLE 2.1. A standard example of a dual pair of crystalline norms is the 1-norm $|\cdot|_1$ and the max-norm $|\cdot|_{\infty}$. For p > 1, the dual norm of a p-norm $|\cdot|_p$ is the q-norm with 1/p+1/q=1. For a positive definite invertible matrix A, the norm $\phi(x) = |Ax|$ has the dual $\phi(x) = |A^{-T}x|$. See Appendix A for general formulas of Wulff shapes for smooth and crystalline ϕ .

Define the (isotropic) total variation of a function $u \in L^1_{loc}(\Omega)$ to be

$$\int_{\Omega} |\nabla u| := \sup\left\{\int_{\Omega} u(x) \operatorname{div} \psi(x) dx : \psi \in C_0^1(\Omega), |\psi(x)| \le 1, \forall x \in \Omega\right\}.$$
(2.5)

The space of all functions $u \in L^1(\Omega)$ such that $\int_{\Omega} |\nabla u| < \infty$ is the set $BV(\Omega)$, for which $\int_{\Omega} |\nabla u|$ becomes a semi-norm.

Generalizing from the Euclidean norm to the norm given by the surface free energy function ϕ , define the *anisotropic total variation* [EO04] as

$$\int_{\Omega} \phi(\nabla u) := \sup\left\{\int_{\Omega} u(x) \operatorname{div} \psi(x) dx : \psi \in C_0^1(\Omega), \phi^{\circ}(\psi(x)) \le 1, \forall x \in \Omega\right\},$$
(2.6)

Write $dist(Y): \Omega \to \mathbb{R}$ for the signed distance function to a set $Y \subset \Omega$, defined by

$$\operatorname{list}(Y)(x) := \inf_{y \in Y} |x - y| - \inf_{y \in \Omega \setminus Y} |x - y|.$$

$$(2.7)$$

Anisotropic Mean Curvature Flow

2.2. Outline of the proposed methods We give a brief outline of the two proposed methods for solving the discrete time evolution of an anisotropic mean curvature flow. For both cases, we need to solve a minimization problem to evolve the mean curvature flow by one discrete time step h > 0. The minimization problem involves the ROF functional (1.3). There are various methods for performing this minimization. We used the split Bregman method which has the advantage that it splits the problem into two simpler problems. The first step is to solve a linear elliptic PDE. The second step is pointwise minimization, which, in some cases, has an explicit solution called *shrinkage*. The additional advantage is that the anisotropy only appears in the second step. In that case we are required to solve an anisotropic shrinkage problem, which we show has an explicit solution in the crystalline case. In the smooth case, no such explicit solution can be obtained. As a result, the methods used to solve this problem are very different in the two cases.

We will now give an outline of the overall method, which will be expanded on in the rest of the article.

The starting point is the isotropic case $\phi(\cdot) = |\cdot|$. The split Bregman method for the ROF functional is described in detail in Section 3.1. It involves alternatingly solving an elliptic PDE (for u^*) and a pointwise minimization,

$$(1/h - \mu\Delta)u^* = f/h - \mu\nabla \cdot (d - b), \tag{2.8}$$

$$d^* = \arg\min_{d} ||d||_{L^1(\Omega)} + \frac{\mu}{2} ||d - \nabla u - b||_{L^2(\Omega)}^2.$$
(2.9)

for a parameter $\mu > 0$. The minimization (2.9) has an explicit solution

$$d^* = \operatorname{shrink}(\nabla u + b, 1/\mu),$$

called *shrinkage*, which is defined for all $x \in \mathbb{R}^2$, $\lambda > 0$ by

shrink
$$(x,\lambda) = \begin{cases} \max(|x| - \lambda, 0) \frac{x}{|x|}, & \text{for } x \neq 0, \\ 0 & \text{for } x = 0. \end{cases}$$

The application of the method to isotropic mean curvature flow is described in Section 3.2.

For a general surface free energy function ϕ , the elliptic solve (2.8) remains unchanged, while the pointwise minimization step (2.9) is replaced by

$$d^* = \arg\min_d \int_\Omega \phi(d) + \frac{\mu}{2} ||d - \nabla u - b||^2_{L^2(\Omega)}$$

The minimization decouples at each point in Ω and thus can be written as

$$d^* = \operatorname{shrink}_{\phi}(\nabla u + b, 1/\mu) \tag{2.10}$$

where

$$\operatorname{shrink}_{\phi}(x,\lambda) := \arg\min_{y \in \mathbb{R}^2} \phi(y) + \frac{\lambda}{2} |x - y|^2.$$
(2.11)

In the case where ϕ is crystalline, we shall call the right hand side of (2.11) a *polyhedral* shrinkage problem. We obtain an explicit formula for the polyhedral shrinkage problem in Section 4.1.

For the case where ϕ is smooth, the method we propose is quite different. No explicit solution for 2.11 is available; instead we consider a system of nonlinear evolution equations which, in the steady state limit, solves both minimizations (2.8) and (2.9) simultaneously. The system is presented in Section 5.1, and derived in Section 5.2.

3. The split Bregman method We first lay out a known algorithm for isotropic mean curvature flow based on the split Bregman method [GBO09]. This will be the prototype for the anisotropic mean curvature flow algorithms described later.

3.1. Overview of the split Bregman method First, we briefly describe the split Bregman method applied to the ROF functional. For further details we refer the reader to [GO09].

Let $\Omega \subset \mathbb{R}^2$ be a bounded, connected set, $f \in BV(\Omega)$, and $\lambda > 0$ be a real parameter. We seek to minimize the ROF functional

$$\int_{\Omega} |\nabla u| + \frac{\lambda}{2} ||u - f||^2_{L^2(\Omega)}$$
(3.1)

over the set of all $u \in L^2(\Omega)$. The key idea is to consider the following constrained minimization problem:

$$\min_{u,d} J(u,d) = ||d||_{L^1(\Omega)} + \frac{\lambda}{2} ||u - f||_{L^2(\Omega)}^2 \quad \text{such that} \quad d = \nabla u.$$
(3.2)

Then, (3.2) is solved by the Bregman iteration [Brè67, OBG+05]:

$$(u^{k+1}, d^{k+1}) = \arg\min_{u, d} D_J^{p_u^k, p_d^k}(u, u^k, d, d^k) + \frac{\mu}{2} ||d - \nabla u||_{L^2(\Omega)}^2$$
(3.3)

$$p_u^k := \partial_u J(u^k, d^k) = \lambda(u^k - f) \tag{3.4}$$

$$p_d^k := \partial_d J(u^k, d^k) = \partial_d ||d^k||_{L^1(\Omega)}$$

$$(3.5)$$

where $\partial_u J(u,k)$ and $\partial_d J(u,k)$ denote the u and d subgradients of J(u,k), respectively, and $D_{J_u^{p_u^k,p_d^k}}^{p_u^k,p_d^k}(u,u^k,d,d^k)$ is the *Bregman distance* of $J(\cdot,\cdot)$, defined by

$$D_{J}^{p_{u}^{k},p_{d}^{k}}(u,u^{k},d,d^{k}) := J(u,d) + J(u^{k},d^{k}) - \left\langle p_{u}^{k},u-u^{k}\right\rangle - \left\langle p_{d}^{k},d-d^{k}\right\rangle.$$
(3.6)

The initial condition is set to $u^0 = f$, $d^0 = p_u^0 = p_d^0 = 0$. Then under suitable conditions, u^k converges to the minimizer of (3.1), see remark 3.1. It can be shown that the iterative sequence (3.3), (3.4) and (3.5) can be equivalently written as,

$$(u^{k+1}, d^{k+1}) = \arg\min_{u, d} ||d||_{L^1(\Omega)} + \frac{\lambda}{2} ||u - f||_{L^2(\Omega)}^2 + \frac{\mu}{2} ||d - \nabla u - b^k||_{L^2(\Omega)}^2$$
(3.7)

$$b^{k+1} = b^k + (\nabla u^{k+1} - d^{k+1}), \tag{3.8}$$

where $b^0 = 0$. The iterative method (3.7) and (3.8) is called the *split Bregman method* for the ROF functional (3.1). To find the minimizer of (3.7), the authors in [GO09] use a two-step alternating minimization:

$$u^{k+1} = \arg\min_{u} \frac{\lambda}{2} ||u - f||^{2}_{L^{2}(\Omega)} + \frac{\mu}{2} ||d^{k} - \nabla u - b^{k}||^{2}_{L^{2}(\Omega)}$$
(3.9)

$$d^{k+1} = \arg\min_{d} ||d||_{L^{1}(\Omega)} + \frac{\mu}{2} ||d - \nabla u^{k+1} - b^{k}||_{L^{2}(\Omega)}^{2}.$$
(3.10)

The first minimization (3.9) can be solved via direct calculus; the Euler-Lagrange equation gives a necessary condition on the minimizing function u^{k+1} :

$$(\lambda - \mu \Delta)u^{k+1} = \lambda f - \mu \nabla \cdot (d^k - b^k)$$
(3.11)

coupled with the boundary condition (see Appendix B for its derivation)

$$\nabla u^{k+1} \cdot n = (d^k - b^k) \cdot n \quad \text{on } \partial\Omega \tag{3.12}$$

where n is the outward normal to $\partial \Omega$.

With respect to the second minimization (3.10), as mentioned in Section 2.2, the problem decouples for every point in Ω . The minimizer is described by the shrinkage operator,

$$\begin{split} d^{k+1} &= \mathrm{shrink}(\nabla u^{k+1} - b^k, 1/\mu) \\ &:= \max(|\nabla u^{k+1} - b^k| - 1/\mu, 0) \frac{\nabla u^{k+1} - b^k}{|\nabla u^{k+1} - b^k|}, \end{split}$$

at each point in Ω .

REMARK 3.1. The split Bregman method utilizes previously known technques in optimization. Setzer [Set09] proved that the split Bregman method converges to the minimizer of (3.1) by establishing that (3.7) and (3.8) is the Augmented Lagragian method (c.f. [NW06, Chaper 17]) and that (3.9) and (3.10) is a special case of the Douglas-Rachford splitting method [EB92].

3.2. Isotropic mean curvature flow We now apply the split Bregman algorithm to Chambolle's formulation for isotropic mean curvature flow.

Consider an initial closed curve $C^0 = \partial S^0$, where S^0 is an open subset in Ω . The goal is to approximate C(t), the mean curvature flow of $C(0) = C^0$, after time t > 0. For a fixed time step h > 0, let $w_{S^0}^* \in L^2(\Omega)$ be the minimizer of the ROF functional (3.1) with $\lambda = 1/h$ and $f = \operatorname{dist}(S^0)$. Define the operator $T_h(S^0) := \{x \in \Omega : w_{S^0}^*(x) < 0\}$. Then, for t > 0, let

$$S_h(t) := (T_h)^{[t/h]}(S^0).$$
(3.13)

It can be shown that $C_h(t) = \partial S_h(t)$ for t > 0 is the discrete time approximation to the mean curvature flow of the initial contour C^0 . Furthermore, (3.13) implements a *monotone* curve evolution of the flow introduced by Almgren, Taylor and Wang [ATW93]; that is, $S \subset S'$ implies $T_h(S) \subset T_h(S')$.

For the actual implementation, it is convenient to work with level set functions rather than sets. Let $u_j(x) := \operatorname{dist}_{(T_h)^j(S^0)}(x)$, for j = 0, 1, 2..., the sequence of distance functions to the sets $S^j := (T_h)^j(S^0)$, for j = 0, 1, 2... Note that the zero level sets of $u_j(x)$ form the discrete time approximation of the evolution of C^0 .

We outline the split Bregman approach to isotropic mean curvature flow in Algorithm 1. The loops in n and m, called the *outer loop* and the *inner loop*, respectively. The inner loop iteratively solves (3.9) and (3.10) to minimize (3.7). The outer loop iteratively performs (3.7) and (3.8). The constants N and M may be chosen manually, or alternatively, the outer and inner loops may be carried out until consecutive iterates are within a given tolerance. The loop in j corresponds to the discrete time iteration of the approximation $C(jh) \approx C_h(jh) = \{u_j(x) = 0\}$.

4. Crystalline Mean Curvature Flow To incorporate the anisotropy of ϕ in the split Bregman formulation of mean curvature flow, one must consider the anisotropic ROF problem:

$$\int_{\Omega} \phi(\nabla u) + \frac{\lambda}{2} ||u - f||_{L^2(\Omega)}^2.$$

$$\tag{4.1}$$

Input: $u_0 = \operatorname{dist}(S^0)$, final time T > 0, parameter $\mu > 0$. **Output**: $u_{[T/h]} = \operatorname{dist}(S_h([T/h])).$ foreach j = 0, 1, ..., [T/h] do Set $f := u_j$; Initialize $u_j^{0,0} := u_j, b^0 = d^{0,0} = 0$; (Outer loop) foreach n = 0, 1, ..., N do (Inner loop) foreach m = 0, 1, ..., M do Solve for $u_j^{n,m+1}$ in $(\frac{1}{h} - \mu \Delta) u_j^{n,m+1} = \frac{1}{h} f - \mu \nabla \cdot (d^{n,m} - b^n)$ with boundary conditions (3.12); Set $d^{n,m+1} := \operatorname{shrink}(\nabla u_j^{n,m+1} - b^n, 1/\mu)$ pointwise in discrete space; end $\begin{array}{l} \text{Set } b^{n+1}\!:=\!b^n\!+\!(\nabla u_j^{n,M}\!-\!d^{n,M});\\ \text{Set } u_j^{n+1,0}\!:=\!u_j^{n,M};\\ \text{Set } d^{n+1,0}\!:=\!d^{n,M}; \end{array}$ end Set $u_{j+1} := dist(\{u_j^{N,M} \le 0\});$

Algorithm 1: Isotropic mean curvature flow using the split Bregman algorithm.

Following the derivation in Section 3.1, the anisotropy appears only in the pointwise minimization (3.10). This leads us to consider the anisotropic shrinkage problem (2.10).

4.1. Polyhedral shrinkage In this section we show that the generalized shrinkage problem is equivalent to the projection onto the Wulff shape. Subsequently, we obtain an explicit solution formula for the generalized shrinkage problem for a *poly*hedral norm (see (4.6) for a definition).

We start by writing ϕ as in (2.4),

$$\phi(x) = \sup_{z \in \mathcal{W}_{\phi}} z \cdot x, \tag{4.2}$$

and define $\pi_{\mathcal{W}_{\phi}}(z)$ to be the projection of z onto \mathcal{W}_{ϕ} ,

$$\pi_{\mathcal{W}_{\phi}}(y) := \arg\min_{x \in \mathcal{W}_{\phi}} |x - y|^2.$$

Consider the quadratic plus norm minimization problem

$$x^* = x^*(y) := \arg\min_{x \in \mathbb{R}^n} \left\{ \phi(x) + \frac{1}{2} |x - y|^2 \right\}.$$
(4.3)

REMARK 4.1. Note the relation of $x^*(y)$ to the generalized shrinkage function introduced in Section 2.2:

$$x^*(y) = shrink_{\phi}(y, 1). \tag{4.4}$$

The anisotropic shrinkage shrink $\phi(y,\lambda)$ for arbitrary $\lambda > 0$ is equivalent to $x^*(y)$, if ϕ in (4.3) is scaled by $1/\lambda$.

Anisotropic Mean Curvature Flow

We claim the following characterization holds. LEMMA 4.1. Denote $x^*(y)$ as the minimizer of (4.3). Then

$$x^{*}(y) = y - \pi_{\mathcal{W}_{\phi}}(y). \tag{4.5}$$

Proof. Rewrite the minimization in (4.3) using (4.2) as

$$\min_{x \in \mathbb{R}^n} \max_{z \in \mathcal{W}_{\phi}} \left\{ \frac{1}{2} |x - y|^2 + z \cdot x \right\}.$$

Since the inner function is convex in x and affine in y, we can interchange the order of min and max to obtain

$$\max_{z \in \mathcal{W}_{\phi}} \min_{x \in \mathbb{R}^n} \left\{ \frac{1}{2} |x - y|^2 + z \cdot x \right\}.$$

Evaluate the inner problem to obtain

$$x^*(y) = y - z,$$

which results in $\max_{z \in \mathcal{W}_{\phi}} \left\{ -\frac{1}{2}z^2 + z \cdot y \right\}$, or equivalently,

$$\min_{z\in\mathcal{W}_{\phi}}\left\{\frac{1}{2}|z-y|^{2}\right\}.$$

The last problem is a projection problem, the minimizer given by

$$z = \pi_{\mathcal{W}_{\phi}}(y).$$

The last result, together with $x^*(y) = y - z$, yields (4.5). PROPOSITION 4.2. $y \in \mathcal{W}_{\phi}$ if and only if $x^*(y) = 0$.

Proof. This is an immediate consequence of (4.5). \Box

From hereon in this Section, we let ϕ be a *polyhedral norm*: given a set of normals $\mathcal{N} = \{n_i\}_{i=1}^k \subset \mathbb{R}^2$ ordered clockwise (define $n_{k+1} = n_1$ and $n_0 = n_k$), write

$$\phi(x) = \max_{n_i \in \mathcal{N}} n_i \cdot x. \tag{4.6}$$

Partition $\mathbb{R}^2 \setminus \{0\}$ into the regions:

$$\begin{split} C_i = & \{ x \neq 0 : n_i \cdot x > n_j \cdot x \text{ for } j = 1, ..., k \} \\ R_{i,i+1} = & \{ x \neq 0 : n_i \cdot x = n_{i+1} \cdot x > 0 \} \\ i = 1, ..., k \end{split}$$

Thus, ϕ is smooth in C_i and has a kink along the rays $R_{i,i+1}$, for each $i=1,\ldots,k$. With Proposition 4.2, we are left with characterizing the minimizer for the case $y \notin W_{\phi}$ (equivalently $\phi^{\circ}(y) > 1$).

THEOREM 4.3 (Polyhedral Shrinkage Formula). Assume that the normals in (4.6) are ordered clockwise, and that $\phi^{\circ}(y) > 1$. Define, for each i = 1, ..., k,

$$\lambda_i = \lambda_i(y) := \frac{(n_i - n_{i+1}) \cdot y + \|n_{i+1}\|^2 - n_i \cdot n_{i+1}}{\|n_i - n_{i+1}\|^2}.$$
(4.7)

Then, there exists a unique $j \in \{1, ..., k\}$ such that either

$$\lambda_j(y) \in [0,1] \tag{4.8}$$

10

or

$$\lambda_j(y) > 1 \text{ and } \lambda_{j-1}(y) < 0. \tag{4.9}$$

Furthermore, in the former case,

$$x^{*}(y) = y - (n_{j}\lambda_{j}(y) + n_{j+1}(1 - \lambda_{j}(y))), \qquad (4.10)$$

and for the latter case,

$$x^*(y) = y - n_j. (4.11)$$

Proof. Start by taking the subdifferential of (4.3),

$$0 \in \partial \phi(x^*) + (x^* - y), \quad x^* = x^*(y). \tag{4.12}$$

The subdifferential $\partial \phi$ can be explicitly solved:

$$\partial \phi(x) = \begin{cases} n_j & \text{if } x \in C_j \text{ for some } j \in \{1, \dots, k\} \\ \operatorname{\mathbf{conv}}\{n_j, n_{j+1}\} & \text{if } x \in R_{j,j+1} \text{ for some } j \in \{1, \dots, k\} \end{cases}$$
(4.13)

In light of Proposition 4.2, $y \notin \mathcal{W}_{\phi}$ implies that $x^*(y) \neq 0$. Since $\{C_i, R_{i,i+1}\}_{i=1}^k$ partitions $\mathbb{R}^2/\{0\}$, it will be sufficient to consider the two cases: either $x^*(y) \in C_j$ or $x^*(y) \in R_{j,j+1}$, for some $j \in \{1, \dots, k\}$.

Rearrange (4.12) and consider the case when $x^*(y) \in R_{j,j+1}$ for some $j \in \{1, ..., k\}$. Then we have

$$y - (n_j \lambda + n_{j+1}(1 - \lambda)) = x^*(y) \tag{4.14}$$

for some unique $\lambda \in [0,1]$. Muliplying (4.14) by n_j and n_{j+1} separately, and noting that $n_j \cdot x^*(y) = n_{j+1} \cdot x^*(y)$, one can solve for λ : the solution is $\lambda = \lambda_j(y)$ as per (4.7). Setting $\lambda = \lambda_j(y)$ in (4.14) gives (4.10).

For the case $x^*(y) \in C_j$ for some $j \in \{1, \dots, k\}$, we have

$$y - n_j = x^*(y).$$
 (4.15)

To see that $\lambda_j(y) > 1$ and $\lambda_{j-1}(y) < 0$ in this case, note that if $x^*(y) \in C_j$,

$$n_j \cdot x^*(y) > n_i \cdot x^*(y), \quad \text{for all } i \neq j.$$

$$(4.16)$$

Since $y = x^*(y) + n_j$, formula (4.7) with the former inequalities imply

$$\begin{split} \lambda_j(y) &= \frac{(n_j - n_{j+1}) \cdot x^*(y)}{\|n_j - n_{j+1}\|^2} + 1 > 1 \\ \lambda_{j-1}(y) &= \frac{(n_{j-1} - n_j) \cdot x^*(y)}{\|n_j - n_{j+1}\|^2} < 0. \end{split}$$

The uniqueness of $x^*(y)$ in the case $\phi^{\circ}(y) > 1$ (by Lemma 4.1) implies the uniqueness of j such that either (4.8) or (4.9) holds. \Box

Theorem 4.3 can be immediately restated as an algorithm for the polyhedral shrinkage problem, see Algorithm 2.

We illustrate the result of the Proposition 4.2 in Figure 4.1, by computing shrinkage for the octagon norm ϕ_8 , where the eight normals are

$$\mathcal{N} = \{ (\pm 1, 0), (0, \pm 1), (\pm 1, \pm 1) / \sqrt{2}, (\pm 1, \pm 1) / \sqrt{2} \}.$$
(4.17)

Algorithm 2: Polyhedral shrinkage algorithm.



Fig. 4.1: Top left: the Frank diagram for the octagon ϕ_8 . Top right: contour plot of $|x^*(y)|$. The dark contour is the zero level set; note that this is the Wulff shape for ϕ_8 . Bottom left and right: a surface plot of the components of $x^*(y) = (x_1(y), x_2(y))$.

4.2. Algorithm for the crystalline mean curvature flow The polyhedral shrinkage algorithm (Algorithm 2) provides the key ingredient to modify Algorithm 1 for crystalline ϕ . We omit presenting the complete algorithm for the crystalline case since it only differs from Algorithm 1 at a single line; namely, within the inner loop, the line "Set $d^{n,m+1} := \operatorname{shrink}(\nabla u_j^{n,m+1} - b^n, 1/\mu)$ pointwise in discrete space" should be replaced by Algorithm 2 with inputs $y = \nabla u_j^{n,m+1} - b^n$ and the set of normals $\{n_i\}_{i=1}^k$ corresponding to the Frank diagram, scaled accordingly by $1/\mu$, see Remark 4.1. REMARK 4.2. As pointed out in [Cha04a], the necessary modification to Algorithm 1 for solving (1.6) with mobility function $\beta(x) = \phi(x)/|x|$ is the last line which should be replaced by:

Set
$$u_{j+1} := \operatorname{dist}_{\phi^{\circ}}(\{u_j^{N,M} \le 0\}),$$

where

$$\operatorname{dist}_{\phi^{\circ}}(Y)(x) := \inf_{y \in Y} \phi^{\circ}(x-y) - \inf_{y \in \Omega \setminus Y} \phi^{\circ}(x-y)$$

$$(4.18)$$

is the signed distance function to a set in the ϕ° metric.

5. Smooth Anisotropic Mean Curvature Flow We saw in the last section that an explicit characterization of shrinkage was possible for a crystalline (or polyhedral) anisotropy on ϕ . However, for smooth anisotropic ϕ , an explicit formula is not available in general. We give an example to illustrate why this formula is not available below.

Instead, the shrinkage formula needs to be computed numerically. One option is to implement an approximation scheme to project onto a smooth unit ball of the dual norm, say by a Newton's method approach, or by means of an efficient distance function solver such as the algorithm proposed in [Tsa02]. Due to the convexity of the unit ball, Newton's method is expected to converge quickly and thus provide accurate and efficient computational solutions to the shrinkage problem. One may also resort to a general Legendre transform solver [Luc97]. In both of the references cited, the computed solution is on a uniform mesh, so interpolation or extrapolation is required.

We shall take a different route, which does not involve any explicit optimization problems. Our approach, motivated by [BGOX06], is to derive an inverse scale space flow arising from the split Bregman method. This approach is slow: each discrete time step of the mean curvature flow is slower than simply solving the equation by the standard level set method, which is the benchmark for speed. While the implementation is not efficient, it complements the method in the crystalline case. We formally derive a nonlinear system of PDE's (5.3) which solves for the minimizer of anisotropic ROF functional (4.1) in the steady state limit. In Section 5.3, we show that this nonlinear flow possesses several interesting properties; in particular, it solves a semi-implicit Euler scheme for the mean curvature flow PDE (1.5).

For an example to illustrate the difficulty of shrinkage for smooth anisotropies, consider the smooth norm

$$\phi(x) = \sqrt{2x_1^2 + x_2^2} , \quad x = (x_1, x_2).$$
(5.1)

Its dual norm $\phi^{\circ}(x) = \sqrt{\frac{1}{2}x_1^2 + x_2^2}$, has a unit ball of an ellipse elongated in the x_1 direction. Thus, shrinkage for ϕ amounts to computing the closest point to this ellipse. A calculus argument shows that such a task requires one to solve a non-trivial quartic equation. One may easily see that deriving a shrinkage formula for a more complicated smooth norm becomes quickly difficult.

Anisotropic Mean Curvature Flow

5.1. Presentation of the inverse scale space equations We begin by presenting the system of PDE's describing the inverse scale space flow in the case of smooth anisotropy. This system arises from (3.3), (3.4) and (3.5), and will be derived in Section 5.2. Throughout this section, we shall decompose the norm ϕ as follows:

$$\phi(x) = \gamma(\omega) ||x||, \quad \omega = \tan^{-1}(x_2/x_1).$$
 (5.2)

The function γ of phase is 2π -periodic and 0-homogeneous in x. Furthermore, we shall assume that ϕ and γ are smooth functions.

The evolution is in the time parameter s, independent of x, which has the role of a parameter. We suppress the parameter x in the notation and write u=u(s), $\theta=\theta(s)$. The inverse scale space equations consist of two differential equations:

$$\frac{\partial\theta}{\partial s} = \frac{p^{\perp} \cdot \nabla u}{\gamma(\theta) + \gamma''(\theta)}, \qquad p = (\cos\theta, \sin\theta),
\frac{\partial u}{\partial s} = h(\Delta u - \nabla \cdot d), \qquad d = (p \cdot \nabla u)p.$$
(5.3)

The initial conditions are

$$\theta(0) = \tan^{-1}(-f_x/f_y),$$

$$u(0) = f.$$
(5.4)

5.2. Formal derivation of the inverse scale space equations In this section, we formally derive the inverse scale space equations (5.3) by taking the Euler-Lagrange equations of the split Bregman functional (3.3).

Consider the Bregman iterative update formulas (3.3), (3.4) and (3.5) for the split Bregman minimization problem (3.2). For clarity, we write out the right hand side of (3.3) completely using the definition of the Bregman distance:

$$\min_{d,u} \int \gamma(\theta) |d| + \frac{1}{2h} \int |u - f|^2 - \left\langle p_u^k, u - u^k \right\rangle - \left\langle p_d^k, d - d^k \right\rangle + \frac{\mu}{2} \int |d - \nabla u|^2, \quad (5.5)$$

where $\theta = \theta(d) = \tan^{-1}(d_2/d_1)$, and $d = (d_1, d_2)$. Recall the subgradients:

$$p_d^k = \partial_{d^k} \int \gamma(\theta(d^k)) |d^k|, \tag{5.6}$$

$$p_u^k = \frac{1}{h}(u^k - f), \quad \text{since } \lambda = 1/h.$$
(5.7)

First, taking the Euler-Lagrange derivative of the functional with respect to d gives,

$$p_d^{k+1} - p_d^k + \mu(d^k - \nabla u^k) = 0.$$
 (5.8)

Observing (5.8) as a forward Euler update scheme over a time variable s with step size μ , it approximates the PDE

$$\frac{\partial p_d}{\partial s} = \nabla u - d. \tag{5.9}$$

For notational simplicity, let us denote d = Rp, for R scalar and $p = (\cos\theta, \sin\theta)$ a unit vector. By the smoothness assumption of γ , for $d \neq 0$,

$$p_{d} = \partial_{d} \int \gamma(\theta) |d| = \frac{d}{|d|} \gamma(\theta) + |d| \gamma'(\theta) \nabla_{d} \theta(d)$$
$$= \frac{d}{|d|} \gamma(\theta) + \frac{d^{\perp}}{|d|} \gamma'(\theta) \quad \text{where } d^{\perp} = (-d_{2}, d_{1})$$
$$= p \gamma(\theta) + p^{\perp} \gamma'(\theta) \quad \text{where } p^{\perp} = (-p_{2}, p_{1}),$$

thus,

$$\begin{aligned} \frac{\partial p_d}{\partial s} &= (-\sin\theta, \cos\theta) \frac{\partial\theta}{\partial s} \gamma(\theta) + (\cos\theta, \sin\theta) \frac{\partial\theta}{\partial s} \gamma'(\theta) \\ &- (\cos\theta, \sin\theta) \frac{\partial\theta}{\partial s} \gamma'(\theta) + (-\sin\theta, \cos\theta) \frac{\partial\theta}{\partial s} \gamma''(\theta) \\ &= p^{\perp} (\gamma(\theta) + \gamma''(\theta)) \frac{\partial\theta}{\partial s}. \end{aligned}$$

Substituting the latter result into (5.9),

$$p^{\perp}(\gamma(\theta) + \gamma''(\theta))\frac{\partial\theta}{\partial s} = \nabla u - Rp.$$
 (5.10)

If we multiply (5.10) by p,

$$\begin{split} & 0 = p \cdot \nabla u - R |p|^2 \\ & R = p \cdot \nabla u. \end{split}$$

If we multiply (5.10) by p^{\perp} ,

$$\begin{split} (\gamma(\theta) + \gamma''(\theta)) \frac{\partial \theta}{\partial s} &= p^{\perp} \cdot \nabla u - Rp^{\perp} \cdot p \\ \frac{\partial \theta}{\partial s} &= \frac{p^{\perp} \cdot \nabla u}{\gamma(\theta) + \gamma''(\theta)}. \end{split}$$

This is the first differential equation in (5.3).

Next taking the derivative of the functional in (5.5) with respect to u,

$$p_u^{k+1} - p_u^k - \mu (\nabla \cdot d^k - \Delta u^k) = 0.$$
 (5.11)

Since the *u* subgradient is explicitly $p_u^k = \frac{1}{h}(u^k - f)$,

$$\frac{1}{h}(u^{k+1}-f) - \frac{1}{h}(u^k-f) - \mu(\nabla \cdot d^k - \Delta u^k) = 0.$$
 (5.12)

Again taking μ to be the step size of a forward Euler discretization, the PDE for u becomes

$$\frac{\partial u}{\partial s} = h(\Delta u - \nabla \cdot d). \tag{5.13}$$

This is the second differential equation in (5.3).

Finally, we derive the initial conditions (5.4). Following [BGOX06], the initial conditions for an inverse scale space flow are d(0) = 0, $p_d(0) = 0$, $p_u(0) = 0$. Immediately, we have R(0) = |d(0)| = 0. Since $0 = p_u(0) = (u(0) - f)$, we also have u(0) = f. Consequently, since $0 = R(0) = p(0) \cdot \nabla u(0) = p(0) \cdot \nabla f$ holds, $p(0) = (-f_y, f_x) \equiv \nabla^{\perp} f$, and therefore, $\theta(0) = \tan^{-1}(-f_x, f_y)$.

5.3. Properties of the inverse scale space flow Next we present analytical properties of the inverse scale space flow (5.3). For what follows, we use the notation from Section 5.2:

$$p = (\cos\theta, \sin\theta), \tag{5.14}$$

$$R = p \cdot \nabla u. \tag{5.15}$$

We prove two main results. The first concerns convergence of ∇u to d as $s \to \infty$. It provides a convergence proof provided that R(s) stays positive in the limit. We have observed that R(s) may be zero for some s > 0 in general but seems to stay positive for all s > 0 for sufficiently smooth convex f.

PROPOSITION 5.1. Assume u, d, θ, R , and p are sufficiently smooth solutions of (5.3) and (5.4). Suppose there exists $s^* > 0$ such that

$$R(s) \ge R_0 > 0, \qquad for \ all \ s > s^*.$$
 (5.16)

Then, $\|\nabla u(s) - d(s)\|_{L^2(\Omega)} \to 0$ as $s \to \infty$.

The second result establishes a connection between the variational formulation and the mean curvature PDE (1.9).

PROPOSITION 5.2. Assume the hypotheses of Proposition 5.1. Further assume that u, d and θ have sufficiently smooth limits as $s \to \infty$, namely u_{∞} , d_{∞} , θ_{∞} , respectively, and $|\nabla u_{\infty}| \neq 0$. Then

$$\frac{u_{\infty} - f}{h} = (\gamma(\theta_{\infty}) + \gamma''(\theta_{\infty}))\nabla \cdot \frac{\nabla u_{\infty}}{|\nabla u_{\infty}|} |\nabla f|.$$
(5.17)

REMARK 5.1. Proposition 5.2 is a consistency result that recovers the level set PDE for mean curvature flow (1.9) from the inverse scale space equations (5.3). That is, solving the inverse scale space equation to steady state is equivalent to a semi-implicit forward Euler scheme for the time dependent PDE (1.9). While explicit forward Euler schemes for elliptic PDE's suffer time step restrictions $h = O(dx^2)$ (where dx is the spacial discretization), the latter property of the inverse scale space flow suggests that it may be free from such restrictions. Note that (5.17) is different from the semiimplicit Euler scheme of [DDE05], in that the curvature term is fully implicit.

We begin by proving several immediate consequences of (5.3) and (5.4): LEMMA 5.3. Assume the hypotheses of Proposition 5.1. Then,

1. $u(s) = h(\gamma(\theta(s)) + \gamma''(\theta(s))) \nabla \cdot p(s) + f$, for all $s \ge 0$,

- 2. $\frac{\partial}{\partial s} \|u(s)\|_{L^2(\Omega)}^2 \leq 0$ (decrease in L^2 energy).
- 3. If $R(s) \ge 0$ then $\frac{\partial}{\partial s} \|\nabla u(s) d(s)\|_{L^2(\Omega)} \le 0$.
- 4. $\frac{\partial \theta}{\partial s}(s) > 0$ for $s \ge 0$ until steady state. (d turns counter-clockwise)

Further assume u, d and θ have sufficiently smooth limits as $s \to \infty$, namely $u_{\infty}, d_{\infty}, \theta_{\infty}$, respectively. Then,

5. $\nabla u_{\infty} = d_{\infty}$.

6. $\frac{\partial}{\partial s}D(u_{\infty}, u(s), d_{\infty}, d(s)) \leq 0.$ (decay of Bregman distance)

Proof. For simplicity of notation, we suppress the inverse scale space parameter s in the proofs. 1. One can verify directly: from the derivation in Section 5.2, $\nabla \cdot p_d = \nabla \cdot (p\gamma + p^{\perp}\gamma) = (\gamma + \gamma'') \nabla \cdot p$. By (5.3), the latter result, and interchanging ∇ and $\partial/\partial s$,

$$\frac{\partial u}{\partial s} = h\nabla \cdot (\nabla u - d) = h\nabla \cdot \frac{\partial p_d}{\partial s} = h\frac{\partial}{\partial s}(\gamma + \gamma'')\nabla \cdot p.$$
(5.18)

Integrating over s, coupled by the initial conditions $u(0) = f, p(0) = \nabla^{\perp} f$, one arrives at the desired result.

2. By direct differentiation,

$$\begin{split} \frac{1}{2} \frac{\partial}{\partial s} \|u\|_{L^{2}(\Omega)}^{2} &= \int_{\Omega} u \frac{\partial u}{\partial s} dx \\ &= h \int_{\Omega} u (\Delta u - \nabla \cdot d) \\ &= h \left(- \int_{\Omega} |\nabla u|^{2} + \int_{\Omega} \nabla u \cdot d \right) \\ &= h \left(- \|\nabla u\|_{L^{2}(\Omega)}^{2} + \|R\|_{L^{2}(\Omega)}^{2} \\ &\leq 0. \end{split}$$

The last inequality holds since by (5.15), $|R| = |p \cdot \nabla u| \le |\nabla u|$.

3. Assume $\gamma+\gamma''>0$ for well-posedness and $R(s)\!\geq\!0.$ Then we can define a seminorm

$$\|\cdot\|_{(\gamma+\gamma'')R} := \langle (\gamma(\theta) + \gamma''(\theta))R(s)\cdot, \cdot \rangle^{1/2}, \qquad (5.19)$$

where $\langle \cdot, \cdot \rangle$ is the L^2 inner product in Ω . Since,

$$\frac{\partial d}{\partial s} = \frac{\partial R}{\partial s} p + R \frac{\partial p}{\partial s} = \frac{\partial R}{\partial s} p + p^{\perp} \frac{\partial \theta}{\partial s}, \qquad (5.20)$$

we have

$$\begin{split} \frac{\partial}{\partial s} \frac{1}{2} \| \nabla u - d \|_{L^{2}(\Omega)}^{2} &= \left\langle \nabla u - d, \frac{\partial}{\partial s} (\nabla u - d) \right\rangle \\ &= -\left\langle \Delta u - \nabla \cdot d, \frac{\partial u}{\partial s} \right\rangle - \left\langle \nabla u - d, \frac{\partial R}{\partial s} p + R p^{\perp} \frac{\partial \theta}{\partial s} \right\rangle \\ &= -\frac{1}{h} \left\| \frac{\partial u}{\partial s} \right\|_{L^{2}(\Omega)}^{2} - \left\langle p \cdot (\nabla u - d), \frac{\partial R}{\partial s} \right\rangle - \left\langle p^{\perp} \cdot (\nabla u - d), R \frac{\partial \theta}{\partial s} \right\rangle \\ &= -\frac{1}{h} \left\| \frac{\partial u}{\partial s} \right\|_{L^{2}(\Omega)}^{2} - \left\langle R - R, \frac{\partial R}{\partial s} \right\rangle - \left\langle (\gamma(\theta) + \gamma''(\theta)) \frac{\partial \theta}{\partial s}, R \frac{\partial \theta}{\partial ts} \right\rangle \\ &= -\frac{1}{h} \left\| \frac{\partial u}{\partial s} \right\|_{L^{2}(\Omega)}^{2} - \left\| \frac{\partial \theta}{\partial s} \right\|_{(\gamma + \gamma'')R}^{2} \\ &\leq 0. \end{split}$$

(5.21)

4. Clearly, $\frac{\partial \theta}{\partial s}(0) = |\nabla^{\perp} f|^2 / (\gamma(\theta(0)) + \gamma''(\theta(0))) > 0$. Consider the first instance s = T for which $\frac{\partial \theta}{\partial s}(T) = 0$. Then by (5.3), $p(T) \cdot \nabla^{\perp} u(T) = 0$, thus $p(T) = \pm \nabla u(T) / |\nabla u(T)|$. Also, by (5.3), $R(T) = \pm |\nabla u|$. Therefore, $d(T) = \nabla u(T)$, and so $\frac{\partial u}{\partial s}(T) = h \nabla \cdot (d(T) - \nabla u(T)) = 0$, i.e. the flow has reached a steady state at s = T.

5. Suppose $\nabla u_{\infty} - d_{\infty} = v$, for some constant $v \in \mathbb{R}^2$. Based on the notation (5.14) and (5.15), denote $p_{\infty} = (\cos \theta_{\infty}, \sin \theta_{\infty})$ and $R_{\infty} = p_{\infty} \cdot \nabla u_{\infty}$. We prove v = 0 by showing that $p_{\infty} \cdot v = p_{\infty}^{\perp} \cdot v = 0$. First,

$$p_{\infty} \cdot v = p_{\infty} \cdot \nabla u_{\infty} - p_{\infty} \cdot d_{\infty}$$
$$= R_{\infty} - R_{\infty} = 0.$$

Next, note that $\frac{\partial \theta_{\infty}}{\partial s} = 0$ by the definition of steady state limit, so

$$p_{\infty}^{\perp} \cdot v = p_{\infty}^{\perp} \cdot \nabla u_{\infty} - p_{\infty}^{\perp} \cdot d_{\infty}$$
$$= (\gamma(\theta_{\infty}) + \gamma''(\theta_{\infty})) \frac{\partial \theta_{\infty}}{\partial s} - p_{\infty}^{\perp} \cdot R_{\infty} p_{\infty}$$
$$= 0 - 0 = 0.$$

Therefore, v = 0.

6. By direct differentiation:

$$\begin{split} \frac{\partial}{\partial s} D(u_{\infty}, u, d_{\infty}, d) &= \left\langle p_d, \frac{\partial d}{\partial s} \right\rangle + \left\langle \frac{u-f}{h}, \frac{\partial u}{\partial s} \right\rangle - \left\langle \frac{1}{h} \frac{\partial u}{\partial s}, u_{\infty} - u \right\rangle \\ &- \left\langle \frac{u-f}{h}, \frac{\partial u}{\partial s} \right\rangle - \left\langle \frac{\partial p_d}{\partial s}, d_{\infty} - d \right\rangle - \left\langle p_d, \frac{\partial d}{\partial s} \right\rangle \\ &= -\left\langle \nabla \cdot (\nabla u - d), u_{\infty} - u \right\rangle - \left\langle \nabla u - d, d_{\infty} - d \right\rangle \\ &= \left\langle \nabla u - d, \nabla u_{\infty} - \nabla u - (d_{\infty} - d) \right\rangle \\ &= - \|\nabla u - d\|_{L^2(\Omega)}^2 \quad \text{since } \nabla u_{\infty} = d_{\infty} \\ &\leq 0. \end{split}$$

Result 4 of Lemma 5.3 shows that the direction of d(s) turns counter-clockwise as s increases. Note that initially, d(0) is perpendicular to $\nabla u(0) = \nabla f$. Thus, d(t)turns counter-clockwise until it coincides with ∇u , at which point $\frac{\partial \theta}{\partial s} = 0$ and steady state is reached. Note that ∇u may also "turn", but does so at a much slower rate, scaled by h.

Result 5 of Lemma 5.3 hinges on the assumption that both u and d converge to steady state solutions. While Proposition 5.1 gives a convergence proof under certain conditions, whether such assumption is generally true is still an open question, although all of our numerical tests confirmed the affirmative.

We close this section by proving the two main propositions.

Proof. [Proof of Proposition 5.1] Again, for simplicity of notation, we suppress the inverse scale space parameter s. We may assume that there exists B > 0 such that,

$$B \ge \gamma(\theta) + \gamma''(\theta) > 0.$$

By (5.10), we can write

$$\|\nabla u - d\|_{L^2(\Omega)}^2 = \left\| (\gamma(\theta) + \gamma''(\theta)) \frac{\partial \theta}{\partial s} \right\|_{L^2(\Omega)}^2.$$

From (5.21) (the steps therein), (5.16) and the last equality, for $s > s^*$,

$$\begin{split} \frac{\partial}{\partial s} \frac{1}{2} \| \nabla u - d \|_{L^{2}(\Omega)}^{2} &\leq - \left\langle \frac{\partial \theta}{\partial s}, \frac{\partial \theta}{\partial s} (\gamma(\theta) + \gamma''(\theta)) R(s) \right\rangle \\ &\leq - \frac{R_{0}}{B} \left\langle \frac{\partial \theta}{\partial s}, \frac{\partial \theta}{\partial s} (\gamma(\theta) + \gamma''(\theta))^{2} \right\rangle \\ &= - \frac{R_{0}}{B} \| \nabla u - d \|_{L^{2}(\Omega)}^{2}. \end{split}$$

By Gronwall's inequality, $\|\nabla u(s) - d(s)\|_{L^2(\Omega)} \le \|\nabla u(0) - d(0)\|_{L^2(\Omega)} e^{-\frac{2R_0}{B}s} = e^{-\frac{2R_0}{B}s}$. so taking $s \to \infty$ we have our result. \Box

Proof. [Proof of Proposition 5.2] Since $\nabla u_{\infty} = d_{\infty}$ (result 5 of Lemma 5.3) and p_{∞} is the unit vector pointing in the direction of d_{∞} , we have

$$p_{\infty} = \frac{\nabla u_{\infty}}{|\nabla u_{\infty}|}.$$
(5.22)

Substituting the above into result 1 of Lemma 5.3 and rearranging gives the desired result. Note that $|\nabla f| = 1$, as f is a signed distance function.

5.4. Inverse scale space algorithm for smooth anisotropy Algorithm 3 outlines the algorithm for smooth anisotropic mean curvature flow using the inverse scale space flow (5.3).

The trivial modification to the inverse scale space method for the Remark 5.2. evolution PDE (1.6) with the mobility function $\beta = \gamma$ is to let the first differential equation in (5.3) be

$$\frac{\partial \theta}{\partial s} = \frac{p^{\perp} \cdot \nabla u}{\gamma(\theta) [\gamma(\theta) + \gamma''(\theta)]},$$

and the preceding results holds true.

Input: $u_0 = \operatorname{dist}(S^0)$, final time T > 0, time step h > 0. **Output**: $u_{[T/h]} = \operatorname{dist}(S_h(T))$. foreach j = 0, 1, ..., [T/h] do Set $f := u_j$; Initialize $\theta(0) := \tan^{-1}(-f_x/f_y), u(0) := f;$ Solve (5.3) until steady state: $u^* := \lim_{s \to \infty} u(s);$ Set $u_{j+1} := dist(\{u^* \le 0\});$

end

Algorithm 3: Anisotropic mean curvature flow using the inverse scale space flow.

6. Numerical Results

6.1. Crystalline mean curvature flow We numerically solved the anisotropic mean curvature flow using the polyhedral shrinkage algorithm (Algorithm 2) within Algorithm 1. The boundary value problem (3.11), (3.12) for u^{k+1} was solved using a standard finite difference discretization on a uniform grid. The distance functions were computed using the fast sweeping method [TCOZ03]; more accurate computations of the signed distance function along the zero level set can be computed using numerical PDE techniques, c.f. [CT08].

The norms (4.6) considered were:

- 1-norm: $\mathcal{N}_1 = \{\pm (1,1), \pm (1,-1)\}.$
- Octagon norm: $\mathcal{N}_8 = \{(\pm 1, 0), (0, \pm 1), (\pm 1, \pm 1)/\sqrt{2}, (\pm 1, \pm 1)/\sqrt{2}\}.$
- A skewed hexagon norm: $\mathcal{N}_{skew} = \{(\pm 1, 0), (0, \pm 1), \pm (1, 1))\}.$ A triangle 'norm': $\mathcal{N}_{tri} = \{(\sqrt{2}, \sqrt{2}), (-1, 0), (0, -1)\}.$

Figure 6.1 shows the results for these anisotropies on a 100×100 grid with h = 0.015. The outer and inner iterations were computed until a consecutive approximations were bounded by 2e-5. Typically, it took less than 20 outer iterations, while the inner iterations ranged from 2 to 500. Notice the emergence of the Wulff shapes as the curves near extinction; see Appendix A for Wulff shapes of anisotropic norms. Note that \mathcal{N}_{tri} does not induce a norm due to its asymmetry, but the algorithm computes its flow nonetheless.



Fig. 6.1: Crystalline mean curvature motion for various anisotropies. Clockwise from top left: \mathcal{N}_1 , \mathcal{N}_8 , \mathcal{N}_{tri} , and \mathcal{N}_{skew}

6.2. Inverse scale space algorithm We first present a sequence of plots showing how the inverse scale space flow (5.3) behaves. For the isotropic case ($\gamma = 1$) with time step h=0.015, Figure 6.2 shows $\nabla u(s)$, d(s) and the contour $\{u=0\}$ at various stages in the iterations over s, where $\Delta s = 0.001$. Note how $\theta(s)$ (the direction of the vector d(s)) moves clockwise as per result 4, Lemma 5.3. The stopping condition was $||u^{n+1} - u^n||_{\infty} \leq 1e-5$, where u^n is the numerical approximation of $u(n\Delta s)$, for which the iteration halted after 3770 iterations. We have observed that the number of iterations necessary for convergence (for the same stopping condition and time step) was roughly between 3000 to 4000 for grid sizes $25^2, 50^2, 100^2$, and 150^2 on the square $[-1,1]^2$. It is reasonable to believe that the number of iterations to convergence should be independent of the grid size, since convergence is attained for a large enough s, which is solely dependent on the underlying equation (i.e. choice of γ , f).

Figure 6.4 shows the plot of $\|\nabla u(s) - d(s)\|_{L^2(\Omega)}$ over iterations in s of (5.3), for



Fig. 6.2: Visualization of ∇u and d in the inverse scale space flow (5.3), with $\gamma_0 = 1$ (the isotropic case). Thick arrows are ∇u and the thin arrows are d. Top to bottom: 200, 1000, 3000 iterations.

various anisotropies. The anisotropies tested were

$$\gamma(\omega) = \gamma_n(\omega) := \frac{1}{n^2 + 1} (n^2 + 1 - \sin(n\omega)), \quad \text{for } n = 0, 2, 4, 8, \tag{6.1}$$

where $\gamma(\omega)$ is defined in (5.2). Note that n=0 is the isotropic case; for n=2,4,8, the anisotropy has a *n*-fold rotational symmetry. We have found that, in general, the isotropic case converges faster to steady state than the anisotropic cases; also, for all cases tested, $\|\nabla u - d\|_{L^2(\Omega)}$ eventually converges exponentially, corroborating with the results of Proposition 5.1.

Results for Algorithm 3 are shown in Figure 6.3 for the anisotropies (6.1).



Fig. 6.3: Mean curvature motion for various smooth anisotropies. Clockwise from top left: γ_0 (isotropic), γ_2 , γ_8 , and γ_4 .

7. Conclusions We have presented new methods of computing crystalline and smooth anisotropic mean curvature flow. In the crystalline case, we proved a formula for the anisotropic shrinkage problem, which was embedded in the split Bregman framework for mean curvature flow. In the smooth anisotropy case, we formally derived an inverse scale space flow, which, in the steady state limit, solves the semi-implicit Euler scheme of the mean curvature PDE. Numerical results for both methods are presented.

Acknowledgement. S. Osher and R. Takei were partially supported by ONR grants N00014-03-1-007 and N00014-07-0810. R. Tsai was partially supported by NSF grant DMS-0714612.

Appendix A. Formulae for Wulff shapes. Assume that ϕ is a convex, positively 1-homogenous function of \mathbb{R}^2 .

The formulas for the Wulff shapes are defined implicitly through the dual norms.



Fig. 6.4: Plot of $\|\nabla u - d\|$ in a log-scale over iterations in the inverse scale space variable s.

However, when we have explicit polyhedral or smooth formulas for the norms, we can also obtain explicit formulas for the Wulff shapes. PROPOSITION A.1. Let ϕ be a polyhedral norm, written explicitly,

$$\phi(x) = \max_{n_i \in \mathcal{N}} n_i \cdot x, \tag{A.1}$$

(\mathcal{N} is the set of normals, as in (4.6)), and let ϕ° be its dual norm. The Wulff shape is the polygon with vertices \mathcal{N} ,

$$\mathcal{W}_{\phi} = \mathbf{conv}(\mathcal{N}).$$

Proof. We show that $\phi^{\circ}(x) = 1$ on the boundary of the polygon $\operatorname{conv}(\mathcal{N})$. The result follows since ϕ° is positively 1-homogeneous. From the definition, for some $n_j \in \mathcal{N}$,

$$\phi^{\circ}(n_{j}) = \max\{n_{j} \cdot y : \max_{n_{j} \in \mathcal{N}} \{n_{i} \cdot y\} = 1\} \le 1$$

By choosing $y = n_j/|n_j|$, we see that $\phi^{\circ}(n_i) = 1$. Next, for $\lambda \in (0,1)$ and two adjacent normals $n_j, n_{j+1} \in \mathcal{N}$,

$$\phi^{\circ}(n_{j}\lambda+n_{j+1}(1-\lambda)) = \max\{n_{j}\cdot y\lambda+n_{j+1}\cdot y(1-\lambda): \max_{n_{i}\in\mathcal{N}}\{n_{i}\cdot y\}=1\}.$$

The maximizing y satisfies $n_j \cdot y = n_{j+1} \cdot y = 1$. Therefore, $\phi^{\circ}(n_j \lambda + n_{j+1}(1-\lambda)) = \lambda + (1-\lambda) = 1$. \Box

Next we describe the Wulff shape for smooth norms.

PROPOSITION A.2. Let ϕ be a smooth norm written explicitly as,

$$\phi(x) = \gamma(\theta_x) \|x\|, \qquad \theta_x = \tan^{-1}(x_2/x_1).$$

Then the Wulff shape is given explicitly as

$$\mathcal{W}_{\phi} = \{ x : \gamma(\theta_x) \ge \|x\| \}.$$

Proof. The dual norm of ϕ is

$$\begin{split} \phi^{\circ}(x) &= \max\{x \cdot y : \gamma(\theta_y) \|y\| = 1\} \quad \text{note: } \theta_y = \tan^{-1}(y_2/y_1) \\ &= \max\{\|x\| \|y\| \cos(\theta_x - \theta_y) : \gamma(\theta_y)\|y\| = 1\} \\ &= \max_{\theta_y} \left\{ \frac{\|x\| \cos(\theta_x - \theta_y)}{\gamma(\theta_y)} \right\}. \end{split}$$

If we write $\phi^{\circ}(x) = \gamma^{\circ}(\theta_x) ||x||$ for the dual norm, we have

$$\gamma^{\circ}(\theta_x) = \max_{\theta_y} \left\{ \frac{\cos(\theta_x - \theta_y)}{\gamma(\theta_y)} \right\}.$$
(A.2)

Assuming $\gamma(\theta_y)$ is smooth, convex and non-vanishing, the maximizer θ_y^* can be characterized by calculus:

$$\tan(\theta_x - \theta_y^*) = \frac{\gamma'(\theta_y^*)}{\gamma(\theta_y^*)}.$$
(A.3)

Furthermore, solving the differential equation for $\gamma(\theta_u^*)$ gives

$$\frac{\cos(\theta_x - \theta_y^*)}{\gamma(\theta_y^*)} = \frac{1}{\gamma(\theta_x)}.$$
(A.4)

Therefore, by (A.3), $\gamma^{\circ}(\theta_x) = 1/\gamma(\theta_x)$; that is, the dual norm is $\phi^{\circ} = ||x||/\gamma(\theta_x)$. The formula for the Wulff shape follows. \Box

Appendix B. Boundary conditions for the Euler-Lagrange equation (3.11). We derive the boundary conditions (3.12), for the minimization

$$\min_{u} I(u) := \frac{\lambda}{2} \int_{\Omega} (u - f)^2 + \frac{\mu}{2} \int_{\Omega} |d^k - \nabla u - b^k|^2,$$
(B.1)

for fixed functions d^k, b^k, f and constants $\lambda, \mu > 0$. To this end, we follow the standard arguments for deriving the first variation, c.f. [Eva10, Chapter 8]: assume u^{k+1} is the minimizer of I(u) and choose any smooth function $v \in C^{\infty}(\Omega)$. Set

$$i(\tau) := I(u^{k+1} + \tau v), \qquad \tau \in \mathbb{R}.$$
(B.2)

By the necessary condition of the minimizer,

$$0 = i'(0) = \lambda \int_{\Omega} (u^{k+1} - f)v + \mu \int_{\Omega} (d^k - \nabla u^{k+1} - b^k)(-\nabla v).$$
(B.3)

Furthermore, taking integration by parts on the second term,

$$0 = \lambda \int_{\Omega} (u^{k+1} - f)v + \mu \int_{\Omega} \nabla \cdot (d^k - \nabla u^{k+1} - b^k)v - \mu \int_{\partial \Omega} (d^k - \nabla u^{k+1} - b^k) \cdot n \ v \ (B.4)$$

where n is the unit outer normal on $\partial\Omega$. The first two terms in (B.4) yields the Euler-Lagrange equation (3.11), while setting the last term to zero gives the desired boundary condition (3.12).

REFERENCES

[ATW93]	Fred Almgren, Jean E. Taylor, and Lihe Wang. Curvature-driven flows: a variational approach. SIAM J. Control Optim., 31(2):387–438, 1993.
[Bel04]	Giovanni Bellettini. A Sampler of Riemann-Finsler Geometry, volume 50 of MSRI Publications, pages 49–82. Campridge University Press, 2004
[BGOX06]	Martin Burger, Guy Gilboa, Stanley Osher, and Jinjun Xu. Nonlinear inverse scale space methods <i>Commun. Math. Sci.</i> 4(1):179–212, 2006
[Bre67]	L. M. Brègman. A relaxation method of finding a common point of convex sets and its application to the solution of problems in convex programming \tilde{Z} Valid Mat i
[DCV00]	Mat. Fiz., 7:620–631, 1967.
[BSV08]	Martin Burger, Christina Stocker, and Axel Voigt. Finite element-based level set meth- ods for higher order flows. J. Sci. Comput., 35(2-3):77–98, 2008.
[CD09]	Antonin Chambolle and Jérôme Darbon. On total variation minimization and surface evolution using parametric maximum flows. Int. J. Comput. Vision, 84(3):288–307, 2009.
[CDK95]	Francine Catté, Françoise Dibos, and Georges Koepfler. A morphological scheme for mean curvature motion and applications to anisotropic diffusion and motion of level acts. SIAM L Numer, Angl. 22(6):1805–1009, 1005
[CGG91]	 Yun Gang Chen, Yoshikazu Giga, and Shun'ichi Goto. Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations. J. Differential Geom., 33(3):749–786, 1991.
[Cha04a]	Antonin Chambolle. An algorithm for mean curvature motion. <i>Interfaces Free Bound.</i> , 6(2):195–218, 2004.
[Cha04b]	Antonin Chambolle. An algorithm for total variation minimization and applications. J. Math. Imaging Vision, 20(1-2):89–97, 2004. Special issue on mathematics and image analysis.
[CHR ⁺ 05]	 Ulrich Clarenz, Frank Haußer, Martin Rumpf, Axel Voigt, and Ulrich Weikard. On level set formulations for anisotropic mean curvature flow and surface diffusion. In Multiscale modeling in epitaxial growth, volume 149 of Internat. Ser. Numer. Math., pages 227–237. Birkhäuser, Basel, 2005.
[CN06]	Antonin Chambolle and Matteo Novaga. Convergence of an algorithm for the anisotropic and crystalline mean curvature flow. <i>SIAM J. Math. Anal.</i> , 37(6):1978–1987, 2006.
[CN07]	Antonin Chambolle and Matteo Novaga. Approximation of the anisotropic mean cur- vature flow. Math. Models Methods Appl. Sci., 17(6):833–844, 2007.
[CT08]	Li-Tien Cheng and Yen-Hsi Tsai. Redistancing by flow of time dependent eikonal equation. J. Comput. Phys., 227(8):4002–4017, 2008.
[DDE05]	Klaus Deckelnick, Gerhard Dziuk, and Charles M. Elliott. Computation of geometric partial differential equations and mean curvature flow. Acta Numer., 14:139–232, 2005.
[DGM96]	Claus Dohmen, Yoshikazu Giga, and Noriko Mizoguchi. Existence of selfsimilar shrink- ing curves for anisotropic curvature flow equations. <i>Calc. Var. Partial Differential Equations</i> , 4(2):103–119, 1996.
[EB92]	Jonathan Eckstein and Dimitri P. Bertsekas. On the Douglas-Rachford splitting method and the proximal point algorithm for maximal monotone operators. <i>Math. Pro-</i> gramming, 55(3, Ser. A):292-318, 1992
[EO04]	Selim Esedoğlu and Stanley J. Osher. Decomposition of images by the anisotropic Budin Ochor Estomi model. <i>Comm. Pure Appl. Math.</i> 57(12):1600–1626–2004
[ERT05]	Selim Esedoglu, Steven Ruuth, and Richard Tsai. Threshold dynamics for shape recon- struction and disocclusion. In <i>In Proc. Int. Conf. Image Processing</i> , pages 502–505, 2005
[ERT08]	Selim Esedoglu, Steven Ruuth, and Richard Tsai. Threshold dynamics for high order
[ERT10]	Selim Esedoğlu, Steven Ruuth, and Richard Tsai. Diffusion generated motion using
[ES91]	L. C. Evans and J. Spruck. Motion of level sets by mean curvature. I. J. Differential
[ET06]	Geom., 53(3):030–081, 1991. Selim Esedoglu and Yen-Hsi Richard Tsai. Threshold dynamics for the piecewise con- stant Mumford-Shah functional <i>L Comput. Phys.</i> 211(1):367–384, 2006
[Eva10]	Lawrence C. Evans. Partial differential equations, volume 19 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, second edition,

	2010
[GBO09]	2010. Tom Goldstein, Xavier Bresson, and Stanley Osher. Geometric applications of the split bregman method: Segmentation and surface reconstruction. Technical report,
[GH08]	UCLA, ftp://ftp.math.ucla.edu/pub/camreport/cam09-06.pdf, July 2009. R. Grzhibovskis and A Heintz. A convolution thresholding scheme for the willmore flow. Interfaces and Free Boundaries 10(2):139-153, 2008
[Gig06]	Yoshikazu Giga. Surface evolution equations, a level set approach, volume 99 of Mono- arrahs in Mathematics. Birkhäuser Varlag. Besel. 2006.
[GL94]	Michael E. Gage and Yi Li. Evolving plane curves by curvature in relative geometries.
[GO09]	Tom Goldstein and Stanley Osher. The split Bregman method for L1-regularized prob- lems SIAM L Imaging Sci. 2(2):323–343, 2009
[Gur93]	Morton E. Gurtin. Thermomechanics of evolving phase boundaries in the plane. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York 1993
[HV06]	Frank Haußer and Axel Voigt. A numerical scheme for regularized anisotropic curve shortening flow Annl Math Lett 19(8):601–698 2006
[LS95]	Stephan Luckhaus and Thomas Sturzenhecker. Implicit time discretization for the mean curvature flow equation. Calc. Var. Partial Differential Equations, 3(2):253–271, 1995
[Luc97]	Yves Lucet. Faster than the fast Legendre transform, the linear-time Legendre trans- form Numer Algorithms 16(2):171–185 (1998) 1997
[MBO94]	Barry Merriman, James K. Bence, and Stanley J. Osher. Motion of multiple functions: a level set approach. J. Commut. Phys. 112(2):334–363, 1994.
[NP07]	Matteo Novaga and Maurizio Paolini. Nonuniqueness for crystalline curvature flow. <i>Math. Models Methods Annl. Sci.</i> 17(8):1307–1315, 2007
[NW06]	Jorge Nocedal and Stephen J. Wright. <i>Numerical optimization</i> . Springer Series in Operations Research and Financial Engineering. Springer, New York, second edition, 2006
[Obe04]	Adam M. Oberman. A convergent monotone difference scheme for motion of level sets
[OBG ⁺ 05]	 Stanley Osher, Martin Burger, Donald Goldfarb, Jinjun Xu, and Wotao Yin. An iterative regularization method for total variation-based image restoration. <i>Multiscale Model. Simul.</i>, 4(2):460–489 (electronic), 2005.
[OS88]	Stanley Osher and James A. Sethian. Fronts propagating with curvature-dependent speed: algorithms based on Hamilton-Jacobi formulations. J. Comput. Phys., 79(1):12–49, 1988.
[RM00]	Steven J. Ruuth and Barry Merriman. Convolution-generated motion and generalized huygens' principles for interface motion. SIAM J. Appl. Math., 60(3):868-890, 2000
[ROF92]	Leoni Rudin, Stanley Osher, and Emad Fatemi. Reconstruction and enhancement of signals using nonlinear non-oscillatory variational methods. <i>Phys. D</i> , 60(1-4):259– 268, 1992
[Ruu98]	Steven J. Ruuth. A diffusion-generated approach to multiphase motion. <i>Journal of Commutational Physics</i> , 145(1):166 – 192, 1998
[Set99]	J. A. Sethian. Level set methods and fast marching methods, volume 3 of Cambridge Monographs on Applied and Computational Mathematics. Cambridge University Press, Cambridge, second edition, 1999. Evolving interfaces in computational ge-
[Set09]	Simon Setzer. Split bregman algorithm, douglas-rachford splitting and frame shrinkage. In XC. Tai, K. Morken, M. Lysaker, and KA. Lie, editors, <i>Scale Space and Variational Methods in Computer Vision</i> , pages 464–476. SSVM, Springer Berlin (Heidelberg, July 2000)
[Tay96]	Jean E. Taylor. Anisotropic interface motion. In Long-Qing Chen, Brent Fultz, John W. Cahn, John R. Manning, John E. Morral, and John Simmons, editors, <i>Mathematics of Microstructure Evolution</i> , volume 4, pages 135–148, SIAM, SIAM, 1996
[TCOZ03]	Yen-Hsi Richard Tsai, Li-Tien Cheng, Stanley Osher, and Hong-Kai Zhao. Fast sweep- ing algorithms for a class of Hamilton-Jacobi equations. <i>SIAM J. Numer. Anal.</i> , 41(2):673–694 (electronic), 2003.
$[T_{co}02]$	Von Hei Richard Tesi. Rapid and accurate computation of the distance function using

[Tsa02] Yen-Hsi Richard Tsai. Rapid and accurate computation of the distance function using grids. Journal of Computational Physics, 178:175–195, 2002.