

# BOUNDARY TREATMENTS FOR MULTILEVEL METHODS ON UNSTRUCTURED MESHES

TONY F. CHAN \*, SUSIE GO \* AND JUN ZOU †

January 24, 1997

**Abstract.** In applying multilevel iterative methods on unstructured meshes, the grid hierarchy can allow general coarse grids whose boundaries may be non-matching to the boundary of the fine grid, so special care is needed to correctly handle different types of boundary conditions. Standard coarse-to-fine grid transfer operators with linear interpolants result in a zero boundary condition for fine grid nodes which are not in the coarse grid domain, and are not accurate enough at Neumann boundaries. In this paper, we propose two effective ways to adapt the standard coarse-to-fine interpolations to correctly implement boundary conditions for two dimensional polygonal domains in such cases: (1) modified coarse grid boundaries and (2) modified interpolations. We prove that all the proposed interpolants possess the local optimal  $L^2$ -approximation and  $H^1$ -stability, which are essential in the convergence analysis of the multilevel Schwarz methods for second order elliptic and parabolic problems on unstructured meshes, and provide some numerical examples of multilevel Schwarz methods (and also multigrid methods) to illustrate the efficiencies of these interpolants.

**Key Words.** Iterative methods, domain decomposition, unstructured meshes

**AMS(MOS) subject classification.** 65F10, 65N30, 65N55

**1. Introduction.** Unstructured grids have become popular in scientific computing because they can be easily adapted to complex geometries and sharp gradients in the solution [3, 12, 17]. However, in order to compete with structured meshes which can exploit the regularity of the mesh, there is a need to develop efficient solvers on unstructured meshes including good multilevel algorithms such as domain decomposition or multigrid methods. Since no natural coarse grids exist as in structured meshes, practical multilevel domain decomposition and multigrid algorithms must allow coarser grids which are non-quasi-uniform, with boundaries and interior elements which are not necessarily matching to that of the fine mesh. The traditional solvers need to be modified so that their efficiency will not be adversely affected by this lack of structure and to ensure that a proper sequence of coarse subspaces exists for the domain decomposition or multigrid methods.

Providing a coarse grid hierarchy for multilevel methods poses some difficulties when using unstructured meshes and several different approaches have been developed recently (see for

---

\* Department of Mathematics, University of California at Los Angeles, Los Angeles, CA 90095-1555, USA. E-mail: chan@math.ucla.edu, sgo@math.ucla.edu. The work was partially supported by ONR under contract ONR-N00014-92-J-1890, and the Army Research Office under contract DAAL-03-91-C-0047 (Univ. Tenn. subcontract ORA4466.04 Amendment 1 and 2). The first two authors also acknowledge support from RIACS under contract number NAS 2-13721 for visits to RIACS/NASA Ames.

† Department of Mathematics, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong. E-mail: zou@math.cuhk.edu.hk. The work of this author was partially supported by the Direct Grant of CUHK and the Hong Kong RGC grant no. CUHK 338/96E.

instance [14, 15, 18, 19]). One technique generates a coarse grid hierarchy by using independent grids created by some grid generator (for example, the one which produced the original grid). Another approach uses agglomeration techniques to create a coarse space hierarchy. Still another method uses a graph approach by forming maximal independent sets (MIS) of the boundaries and interiors of the mesh and then retriangulating the resulting vertex set. The advantage of using a maximal independent set approach is that the grids are node-nested and thus efficient methods can be used to create the interpolation and restriction operators needed to transfer information from one level to the other. A disadvantage however, is that for complicated geometries, particularly in three dimensions, special care must be taken to ensure that the coarse grids which are produced are valid and preserve the important geometric features of the fine domain.

It was shown in [7] that for domain decomposition methods for elliptic problems on unstructured meshes, the same optimal convergence rate can be achieved as in the structured case provided that the coarse grid domain covers the Neumann boundary part of the fine grid domain, but no such requirement is needed for homogeneous Dirichlet boundary conditions. This was demonstrated numerically in [7] with problems on the unit square by physically extending the coarse grid domain beyond the Neumann boundaries and using linear interpolation.

In this paper, we will extend this idea to include interpolants with non-zero extensions which do not require the coarse grid domain be modified to cover the Neumann boundary part of the fine grid domain and provide some analysis on a crucial step in the convergence analysis of multilevel Schwarz methods on unstructured meshes using such coarse-to-fine interpolants. We will follow the general framework for convergence analyses applicable to unstructured meshes in [7, 8, 9], which can be viewed as a natural extension of the one formulated by Xu [23] for structured meshes. Some preliminary results can be found in [5].

This paper is arranged as follows: The considered elliptic problem is introduced in Section 2 and the coarse-to-fine grid transfer operators along with several particular interpolants are defined in Section 3. Section 4 demonstrates the optimal  $L^2$ -approximation and  $H^1$ -stability properties of the interpolants. In Section 5, we provide some numerical results on multilevel Schwarz (cf. [2] [24]) and multigrid methods using the coarse-to-fine grid transfer operators proposed in Section 3. Previous numerical results on multilevel Schwarz methods on structured grids can be found in [21, 24]. The results we present here, however, appear to be the first on multilevel Schwarz on unstructured grids. We summarize some conclusions in Section 6.

**2. The elliptic problem.** Let us consider the following boundary value problem:

$$\begin{aligned} Lu &= f & \text{in } \Omega, \\ u &= 0 & \text{on } \Gamma_D, \\ \frac{\partial u}{\partial n} &= 0 & \text{on } \Gamma_N, \end{aligned}$$

where  $L$  is a second order self-adjoint and uniformly elliptic operator,  $\Omega$  a polygonal domain,  $\Gamma_D$  and  $\Gamma_N$  two curves consisting of piecewise straight lines, with  $\bar{\Gamma}_D \cup \bar{\Gamma}_N = \partial\Omega$ .

We discretize the problem using piecewise linear finite elements and solve the resulting system of equations with the preconditioned conjugate gradient or GMRES method. Let  $\mathcal{T}^h$  be a given fine triangulation of the domain  $\Omega$  with triangular elements, and  $V^h$  be the piecewise linear finite element space defined on  $\mathcal{T}^h$ . Suppose  $\mathcal{T}^H$  is a coarse triangulation of the domain  $\Omega$ , with its elements forming a polygonal domain  $\Omega^H$ . With unstructured meshes, the MIS coarsening strategy for generating a coarse grid hierarchy may produce coarse grid domains whose boundaries do not match that of the fine domain. Note then that  $\Omega^H$  is allowed to be non-nested and non-matching with  $\Omega$ , so in general we have  $\Omega^H \neq \Omega$  (see Fig. 1). Moreover, we do not require the coarse grid  $\mathcal{T}^H$  to have anything to do with the fine grid  $\mathcal{T}^h$ , i.e. none

of the nodes of  $\mathcal{T}^H$  need be nodes of  $\mathcal{T}^h$ , but only that it is shape regular. No assumption on quasi-uniformity is made on the grids  $\mathcal{T}^h$  and  $\mathcal{T}^H$ . Let  $V^H$  be the piecewise linear finite element space corresponding to the coarse grid triangulation  $\mathcal{T}^H$  and the boundary condition in  $V^H$  be defined as follows: each boundary node  $x_i^H \in \partial\Omega^H$  in  $\mathcal{T}^H$  is assigned the same boundary condition type (Dirichlet or Neumann) as the closest fine boundary node to  $x_i^H$ . By changing boundary conditions for a few coarse boundary nodes, if needed, we may assume the coarse boundary nodes are arranged in the way that two neighbors of each Neumann (resp. Dirichlet) node are also of Neumann (resp. Dirichlet) type, with only two Neumann (resp. Dirichlet) nodes near two junctions between  $\Gamma_D$  and  $\Gamma_N$  to have one Dirichlet and one Neumann node as its two neighbors.

It is intuitively obvious that for the coarse grid,  $\mathcal{T}^H$ , to assist in accelerating the convergence of iterative methods on the fine grid,  $\mathcal{T}^h$ , it cannot be allowed to be too small compared with the fine grid. Therefore, we always assume that  $\Omega^H$  covers a significant part of  $\Omega$ . More accurately, we assume that there exists a positive constant  $C$  such that for any point  $x \in \partial\Omega$ , we have

$$\text{dist}(x, \tau^H) \leq C d(\tau^H),$$

where  $\tau^H$  is the closest element in  $\mathcal{T}^H$  to  $x$  and  $d(\tau^H)$  the diameter of  $\tau^H$ .

**3. Coarse-to-fine interpolations.** As the coarse grid boundary  $\partial\Omega^H$  does not match with the the original boundary  $\partial\Omega$ , so the coarse space  $V^H$  is usually not a subspace of the fine space  $V^h$ . In fact, even if  $\Omega^H = \Omega$ ,  $V^H$  may still not be a subspace of  $V^h$  as the coarse elements are often not the unions of some fine elements in the unstructured grid. To construct a coarse-to-fine transfer operator, one may easily come up with the standard nodal value interpolant associated with the fine space  $V^h$ . But notice that this interpolant is only well defined for those fine nodes lying also in the coarse domain  $\bar{\Omega}^H$ , and meaningless for those fine nodes lying outside  $\bar{\Omega}^H$ . A simple and natural way to remove this barrier is to assign those fine node values by zero. This is indeed a reasonable and efficient thing to do when the assignment is done along the coarse boundary part of Dirichlet type (which is also near the fine boundary part of Dirichlet type). We shall denote this interpolant as the coarse-to-fine interpolant,  $\mathcal{I}_h^0$ .

$\mathcal{I}_h^0$ : **Zero extension with unmodified coarse boundaries.** Where coarse grid boundary conditions are of Dirichlet type, the standard nodal value interpolants with zero extensions can be accurate enough for interpolating fine grid values outside the coarse grid domain  $\Omega^H$  (cf. Fig. 1a), we refer to [6, 7] for the theoretical and numerical justifications of  $\mathcal{I}_h^0$ .

Although the interpolant  $\mathcal{I}_h^0$  is appropriate to use at Dirichlet boundaries it is not accurate enough, or not accurate at all sometimes, to use at Neumann boundaries, see the numerical results in [7] and Section 5. To achieve better efficiency, we should modify this intergrid operator to account for the Neumann condition. We now propose two general ways to treat such boundaries:

1. Modify the coarse grid domain to cover any fine grid boundaries of Neumann type.
2. Increase the accuracy of the interpolants by accounting for the Neumann condition for those fine nodes in  $\Omega \setminus \Omega^H$ .

The first approach is motivated by the fact that standard nodal value interpolants can still be used with efficiency where the coarse grid covers the Neumann boundary part of the fine grid. This was first proposed and justified in [7]. We shall denote this operator as the coarse-to-fine interpolant,  $\mathcal{I}_h^1$ .

$\mathcal{I}_h^1$ : **Zero extension with modified coarse boundaries.** Modify the original coarse grid domain  $\Omega^H$  to make it appropriately larger so that it covers the Neumann boundary part of the fine grid domain (see Fig. 1b). Let us still denote the modified coarse grid domain by  $\Omega^H$ .

Then for all  $v^H \in V^H$ , the interpolant  $\mathcal{I}_h^1$  is defined as

$$\mathcal{I}_h^1 v^H(x_j^h) = \begin{cases} v^H(x_j^h) & \text{for } x_j^h \in \Omega \cap \bar{\Omega}^H, \\ 0 & \text{for } x_j^h \in \Omega \setminus \bar{\Omega}^H. \end{cases}$$

This is a natural extension of  $v^H$  by zero outside the Dirichlet boundary part of the coarse grid domain. Similar zero extensions were used in Kornhuber-Yserentant [16] to embed an arbitrarily complicated domain into a square or cube in constructing multilevel methods on nested and quasi-uniform meshes for second order elliptic problems with purely Dirichlet boundary conditions.




FIG. 1. *Zero extension interpolants: a)  $\mathcal{I}_h^0$ : With unmodified coarse boundaries; b)  $\mathcal{I}_h^1$ : With modified coarse boundaries to cover the parts where Neumann conditions exist (dashed lines). Thick lines represent coarse grid boundaries.*

Although the coarse-to-fine operator  $I_h^1$  works well for mixed boundary conditions, one has to modify the original coarse grid so that it covers the Neumann boundary part of the fine grid domain. This can be difficult to do for very complicated domains. To avoid modifying the original coarse grid, we now consider standard finite element interpolants which are modified only near Neumann boundaries. To do so, we first introduce some notation. Let  $\tau_{lr}^H$  be any coarse boundary element in  $\mathcal{T}^H$  made up of the three vertices  $x_l^H, x_r^H, x_i^H$  and which has an edge on the boundary  $\partial\Omega^H$ , denoted by  $x_l^H x_r^H$ . We use  $\Omega(x_l^H, x_r^H)$  to denote the union of all fine elements, if any, which has a non-empty intersection with the unbounded domain formed by the edge  $x_l^H x_r^H$  and two outward normal lines to  $x_l^H x_r^H$  at two vertices  $x_l^H, x_r^H$  (cf. Fig. 2). By including a few more fine elements in some  $\Omega(x_l^H, x_r^H)$ , if necessary, we may assume that the fine grid part ( $\Omega \setminus \Omega^H$ ) is included in the union of all  $\Omega(x_l^H, x_r^H)$ . Moreover, we assume

$$(H1) \quad \text{diam } \Omega(x_l^H, x_r^H) \leq \mu_0 \text{diam } \tau_{lr}^H$$

which implies the measure of  $\Omega(x_l^H, x_r^H)$  is bounded by the measure of  $\tau_{lr}^H$ :

$$|\Omega(x_l^H, x_r^H)| \leq \mu |\tau_{lr}^H|,$$

where  $\mu_0$  and  $\mu$  are two positive constants independent of  $H$  and  $h$ . Without any difficulty, the constant  $\mu_0$ , and so  $\mu$ , can be allowed in our subsequent results to depend on the two nodes  $x_l^H, x_r^H$ . In this case,  $\mu_0$  and  $\mu$  will enter all the related bounds naturally.

We remark that (H1) restricts the size of the fine grid part near the edge  $x_l^H x_r^H$  but outside the coarse grid domain  $\Omega^H$ , that is, each local fine grid part  $\Omega(x_l^H, x_r^H)$  is not allowed to be too large compared to its nearest coarse element  $\tau_{lr}^H$ . This is a reasonable requirement in applications.

Then the standard nodal value interpolant associated with the fine space  $V^h$  can be generalized outward to each local fine grid part  $\Omega(x_l^H, x_r^H)$  using three given linear functions  $\theta_1, \theta_2$  and  $\theta_3$  which are defined in  $\Omega \cup \bar{\Omega}^H$  but bounded in  $\Omega(x_l^H, x_r^H) \cup \tau_{lr}^H$  and satisfy

$$(3.1) \quad \theta_1(x) + \theta_2(x) + \theta_3(x) = 1, \quad \forall x \in \bar{\Omega} \cup \bar{\Omega}^H.$$




FIG. 2. Shaded region,  $\Omega(x_i^H, x_r^H)$ , shows the fine grid part which is not completely covered by the coarse grid domain.

Note that the functions  $\theta_1$ ,  $\theta_2$  and  $\theta_3$  above are not necessarily non-negative, and though they are element  $\tau_{lr}^H$ -related, we will not use any index to specify this relation in order to simplify the notation. Then for any coarse function  $v^H \in V^H$ , we define an operator  $\Theta_h$  by

$$\Theta_h v^H(x) = \theta_1(x)v^H(x_i^H) + \theta_2(x)v^H(x_r^H) + \theta_3(x)v^H(x_i^H), \quad \forall x \in \Omega(x_i^H, x_r^H) \cup \tau_{lr}^H,$$

and assume that

$$(H2) \quad \Theta_h v^H = v^H \quad \text{on the edge } x_i^H x_r^H,$$

which means  $\Theta_h v^H$  is indeed an extension of  $v^H$ . For convenience, later on we will always regard  $\Theta_h v^H$  as a function defined also outside  $\Omega(x_i^H, x_r^H) \cup \tau_{lr}^H$  by extending it naturally.

With the above notation, we can introduce the general coarse-to-fine interpolant  $\mathcal{I}_h$ :

DEFINITION 3.1. For any coarse function  $v^H$  in  $V^H$ , its image under the coarse-to-fine interpolant  $\mathcal{I}_h$  is specified as follows:

(C1) For any fine node  $x_j^h$  in  $\bar{\Omega} \cap \bar{\Omega}^H$ ,

$$\mathcal{I}_h v^H(x_j^h) = v^H(x_j^h);$$

(C2) For any fine node  $x_j^h$  in  $\Omega(x_i^H, x_r^H) \setminus \bar{\Omega}^H$  with both  $x_i^H$  and  $x_r^H$  of Neumann nodes,

$$\mathcal{I}_h v^H(x_j^h) = \Theta_h v^H(x_j^h);$$

(C3) For any fine node  $x_j^h$  in  $\Omega(x_i^H, x_r^H) \setminus \bar{\Omega}^H$  with both  $x_i^H$  and  $x_r^H$  of Dirichlet nodes,

$$\mathcal{I}_h v^H(x_j^h) = 0;$$

(C4) For any fine node  $x_j^h$  in  $\Omega(x_i^H, x_r^H) \setminus \bar{\Omega}^H$  with one of  $x_i^H$  and  $x_r^H$  the Neumann node and one the Dirichlet node,

$$\begin{aligned} \mathcal{I}_h v^H(x_j^h) &= 0, \quad \text{if } x_j^h \text{ is a fine boundary node of Dirichlet type;} \\ \mathcal{I}_h v^H(x_j^h) &= \Theta_h v^H(x_j^h), \quad \text{otherwise.} \end{aligned}$$

The following are two concrete examples of interpolants which satisfy the above definition and assumptions. We only give the corresponding forms of  $\Theta_h$ 's required in the definition:

$\mathcal{I}_h^2$ : **Nearest edge interpolation.** Define the interpolant at  $x_j^h$  by using the nodes of the coarse boundary edge closest to  $x_j^h$  (see Fig. 3):

$$\mathcal{I}_h^2 v^H(x_j^h) = \lambda(x_j^h)v^H(x_i^H) + (1 - \lambda(x_j^h))v^H(x_r^H),$$




FIG. 3. *More accurate interpolants: a)  $\mathcal{I}_h^2$ : Fine nodal values outside the coarse domain are interpolated with coarse nodal values on the nearest coarse grid edge; b)  $\mathcal{I}_h^3$ : Fine nodal values outside the coarse domain are interpolated with nodal values on the nearest coarse element  $\tau_{lr}^H$ . Thick lines represent coarse grid boundaries or elements, and dotted lines show the coarse nodes used to interpolate the fine nodal value at  $x_j^h$ .*

where  $x_l^H$  and  $x_r^H$  are the nodes of the coarse boundary edge closest to  $x_j^h$ , and  $\lambda$  is the ratio of the lengths of two segments of the edge  $x_l^H x_r^H$  cut off by the normal line passing through  $x_j^h$  to the edge (see Fig. 3). This kind of interpolation was used by Bank and Xu [1] in their construction of a hierarchical basis on an unstructured mesh.

$\mathcal{I}_h^3$ : **Nearest element interpolation.** Define the non-zero extension by using barycentric functions (see Fig. 3):

$$\mathcal{I}_h^3 v^H(x_j^h) = \lambda_l(x_j^h) v^H(x_l^H) + \lambda_r(x_j^h) v^H(x_r^H) + \lambda_i(x_j^h) v^H(x_i^H),$$

where  $\lambda_l, \lambda_r, \lambda_i$  are three barycentric coordinate functions (also known as area or volume coordinates) corresponding to  $\tau_{lr}^H$ .

**Remark 3.4.** Note that the functions  $\lambda_l, \lambda_r$  and  $\lambda_i$  used in the definition of  $\mathcal{I}_h^3$  satisfies  $\lambda_l, \lambda_r, \lambda_i \geq 0$  for  $x_j^h \in \tau_{lr}^H$ , but not so for  $x_j^h \notin \tau_{lr}^H$ . In the case as shown in Figure 3b), we have  $x_j^h \notin \tau_{lr}^H$ ,  $\lambda_l(x) \geq 0, \lambda_r(x) \geq 0$ , but  $\lambda_i(x) \leq 0$  and  $\lambda_l(x) + \lambda_r(x) + \lambda_i(x) = 1$ . By (H1), we always have

$$|\lambda_l(x)| \leq \mu_1, \quad |\lambda_r(x)| \leq \mu_1, \quad \text{and} \quad |\lambda_i(x)| \leq \mu_1, \quad \forall x \in \Omega(x_l^H, x_r^H) \cup \tau_{lr}^H,$$

where  $\mu_1$  is a constant independent of  $h$  and  $H$  but depending only on the constant  $\mu$  in (H1).

**4. Stability and approximation properties of the interpolation operator.** The convergence proof for the overlapping multilevel domain decomposition and multigrid methods require the coarse-to-fine grid transfer operator to possess the local optimal  $L^2$ -approximation and local  $H^1$ -stability properties [7, 8, 9]. The locality of these properties is essential to the effectiveness of these methods on highly non-quasi-uniform unstructured meshes.

Purely for our theoretical analysis, we now introduce a triangulation  $\tilde{\mathcal{T}}^H$ . Extend  $\mathcal{T}^H$  to a larger but still shape regular triangulation  $\tilde{\mathcal{T}}^H$ , the corresponding domain denoted by  $\tilde{\Omega}^H$ , such that the Neumann boundary of  $\Omega^H$  is contained in  $\tilde{\Omega}^H$  but the Dirichlet boundary remains the same. Let  $\tilde{V}^H$  be the corresponding piecewise linear finite element space on  $\tilde{\mathcal{T}}^H$  with completely homogeneous Dirichlet boundary condition. Then we have

$$V^H = \tilde{V}^H|_{\Omega^H}.$$

For simplicity, for any coarse element  $\tau^H$  in  $\mathcal{T}^H$ , we let

$$d(\tau^H) = \text{diam}(\tau^H), \quad N(\tau^H) = \text{union of coarse elements adjacent to } \tau^H.$$

Similarly,  $d(\tau^h)$  and  $N(\tau^h)$  are defined for any fine element  $\tau^h$  in  $\mathcal{T}^h$ . We then have the following local optimal  $L^2$ -approximation and  $H^1$ -stability for the operator  $\mathcal{I}_h$  on the coarse space  $V^H$ .

LEMMA 4.1. *Let  $\mathcal{I}_h$  be any interpolation operator defined in Definition 3.1 and  $v^H$  any coarse function in  $V^H$ , if we extend  $v^H$  onto  $\tilde{V}^H$  in any way, still denoted by  $v^H$ , then for any  $\tau^H \in \mathcal{T}^H$ , we have*

$$\begin{aligned}
(1) \quad & \sum_{\substack{\tau^h \cap \tau^H \neq \emptyset \\ \tau^h \subset \bar{\Omega}^H}} \|v^H - \mathcal{I}_h v^H\|_{0,\tau^h}^2 \leq C d^2(\tau^H) |v^H|_{1,N(\tau^H)}^2, \\
(2) \quad & \sum_{\substack{\tau^h \cap \tau^H \neq \emptyset \\ \tau^h \subset \bar{\Omega}^H}} |\mathcal{I}_h v^H|_{1,\tau^h} \leq C |v^H|_{1,N(\tau^H)}, \\
(3) \quad & \sum_{\tau^h \in \Omega(x_i^H, x_r^H)} \|v^H - \mathcal{I}_h v^H\|_{0,\tau^h}^2 \leq C d^2(\tau_r^H) \sum_{\tau^H \in N(\tau_r^H)} |v^H|_{1,N(\tau^H)}^2, \\
(4) \quad & \sum_{\tau^h \in \Omega(x_i^H, x_r^H)} |\mathcal{I}_h v^H|_{1,\tau^h}^2 \leq C \sum_{\tau^H \in N(\tau_r^H)} |v^H|_{1,N(\tau^H)}^2,
\end{aligned}$$

where  $\Omega(x_i^H, x_r^H)$  is any region as introduced in Section 3.

*Proof.* The inequalities (1) and (2) correspond to the parts where the fine grid domain is completely contained in the coarse grid domain. Their proofs can be found in [7, 8]. The last two inequalities (3) and (4) correspond to the fine grid parts which are not covered by the coarse grid and which we shall prove here. We give the proofs only for the cases (C1) – (C2), the other case can be proved similarly.

We first prove inequality (3), i.e.  $L^2$ -optimal approximation. For any fine element  $\tau^h$  in  $\Omega(x_i^H, x_r^H)$ , as  $\mathcal{I}_h v^H$  is linear on  $\tau^h$  we can express

$$\mathcal{I}_h v^H(x) = \sum_{i=1}^3 \mathcal{I}_h v^H(x_i^h) \phi_i^h,$$

where  $x_i^h$  ( $i = 1, 2, 3$ ) are the three vertices of  $\tau^h$  and  $\phi_i^h$  ( $i = 1, 2, 3$ ) the corresponding basis functions of  $V^h$  at these three nodes. Then by definition of  $\mathcal{I}_h$  and the boundedness of  $\theta_i$  ( $i = 1, 2, 3$ ) we have

$$\begin{aligned}
\|\mathcal{I}_h v^H\|_{0,\tau^h}^2 & \leq C d^2(\tau^h) \sum_{i=1}^3 (\mathcal{I}_h v^H(x_i^h))^2 \\
& \leq C d^2(\tau^h) \left\{ (v^H(x_i^H))^2 + (v^H(x_r^H))^2 + (v^H(x_i^H))^2 \right\}.
\end{aligned}$$

Summing over all  $\tau^h \in \Omega(x_i^H, x_r^H)$  and using (H1),

$$\begin{aligned}
\sum_{\tau^h \in \Omega(x_i^H, x_r^H)} \|\mathcal{I}_h v^H\|_{0,\tau^h}^2 & \leq C \left\{ (v^H(x_i^H))^2 + (v^H(x_r^H))^2 + (v^H(x_i^H))^2 \right\} \sum_{\tau^h} d^2(\tau^h) \\
& \leq C \left\{ (v^H(x_i^H))^2 + (v^H(x_r^H))^2 + (v^H(x_i^H))^2 \right\} |\tau_r^H| \\
& \leq C \|v^H\|_{0,\tau_r^H}^2.
\end{aligned}$$

Using this and inequality  $(a+b)^2 \leq 2(a^2+b^2)$ ,  $\forall a, b \in R^1$ , we obtain

$$\begin{aligned}
\sum_{\tau^h \in \Omega(x_i^H, x_r^H)} \|v^H - \mathcal{I}_h v^H\|_{0,\tau^h}^2 & \leq 2 \sum_{\tau^h \in \Omega(x_i^H, x_r^H)} \|v^H\|_{0,\tau^h}^2 + C \|v^H\|_{0,\tau_r^H}^2 \\
& \leq C \|v^H\|_{0,N(\tau_r^H)}^2.
\end{aligned}$$

Noting that the left-hand side of the inequality doesn't change by replacing  $v^H$  by  $v^H$  plus any constant, we obtain by Poincaré inequality that

$$\sum_{\tau^h \in \Omega(x_i^H, x_r^H)} \|v^H - \mathcal{I}_h v^H\|_{0, \tau^h}^2 \leq C d^2(\tau_{lr}^H) |v^H|_{1, N(\tau_{lr}^H)}^2.$$

This proves (13).

We next prove (14), i.e.  $H^1$ -stability. For the ease of notation, we assume that  $\Omega(x_i^H, x_r^H)$  can be covered by  $\tau_{lr}^H$  plus two extended coarse elements  $\tau_1^H$  and  $\tau_2^H \in \tilde{\Omega}^H$  (see Fig. 4).




FIG. 4. Fine grid element (shaded) in  $\Omega(x_i^H, x_r^H)$  which is covered by  $\tau_{lr}^H$  plus two extended coarse elements,  $\tau_1^H$  and  $\tau_2^H$ .

Let us define

$$\tilde{\Theta}_h v^H(x) = \begin{cases} v^H(x), & \text{if } x \in \tilde{\Omega} \cap \tilde{\Omega}^H \\ \Theta_h v^H(x), & \text{if } x \in \tilde{\Omega} \setminus \tilde{\Omega}^H. \end{cases}$$

It is easy to see that  $\tilde{\Theta}_h v^H$  is continuous and belongs to  $H^1(\tilde{\tau}_{lr}^H \cup \Omega(x_i^H, x_r^H))$ , and by definition of  $\mathcal{I}_h$  and  $\tilde{\Theta}_h$ , we have

$$\mathcal{I}_h v^H(x) = \tilde{\mathcal{I}}_h \tilde{\Theta}_h v^H(x), \quad \forall x \in \tilde{\tau}_{lr}^H \cup \Omega(x_i^H, x_r^H).$$

Here  $\tilde{\mathcal{I}}_h$  is the standard nodal value interpolant defined on the finite element space  $V^h$ . We have to bound  $|\mathcal{I}_h v^H|_{1, \tau^h}$  for all  $\tau^h \in \Omega(x_i^H, x_r^H)$ . By the triangle inequality,

$$(4.2) \quad |\mathcal{I}_h v^H|_{1, \tau^h}^2 \leq 2|\tilde{\mathcal{I}}_h \tilde{\Theta}_h v^H - \tilde{\Theta}_h v^H|_{1, \tau^h}^2 + 2|\tilde{\Theta}_h v^H|_{1, \tau^h}^2.$$

For the first term in (4.2), we have by standard interpolation theory (see Ciarlet [10]),

$$(4.3) \quad (I)_3 \equiv |\tilde{\mathcal{I}}_h \tilde{\Theta}_h v^H - \tilde{\Theta}_h v^H|_{1, \tau^h}^2 \leq C h^2 |\tilde{\Theta}_h v^H|_{1, \infty, \tau^h}^2.$$

Let the maximum of  $\tilde{\Theta}_h v^H$  be reached at some point  $x_0$  which must belong to either  $\tau_1^H \cup \tau_2^H$  or  $\tau_{lr}^H$  or  $N(\tau_{lr}^H) \setminus \tau_{lr}^H$ , and denote it by  $m(x_0) = |\tilde{\Theta}_h v^H|_{1, \infty, \tau^h}$ . We consider only the two cases:  $x_0 \in \tau_1^H \cup \tau_2^H$  or  $x_0 \in \tau_{lr}^H$  as the case of  $x_0 \in N(\tau_{lr}^H) \setminus \tau_{lr}^H$  is similar to the one for  $x_0 \in \tau_{lr}^H$ . For either case, we can always construct a shape regular element  $\tau_1^h$  with  $x_0$  as one of its vertices such that  $\tau_1^h \subset \tau_1^H \cup \tau_2^H$  for the former and  $\tau_1^h \subset \tau_{lr}^H$  for the latter and  $d(\tau_1^h)$  is of the same size as  $d(\tau^h)$  (see Fig. 5). Then it follows from the inverse inequality that for  $x_0 \in \tau_1^H \cup \tau_2^H$ ,

$$(I)_3 \leq C d^2(\tau^h) m(x_0) \leq C d^2(\tau^h) |\tilde{\Theta}_h v^H|_{1, \infty, \tau_1^h}^2 \leq C |\tilde{\Theta}_h v^H|_{1, \tau_1^h}^2;$$






FIG. 5. A shape regular element,  $\tau_1^h$ , whose diameter is of the same size as  $\tau^h \in \Omega(x_l^H, x_r^H)$  (shaded).

while for  $x_0 \in \tau_{lr}^H$ ,

$$(I)_3 \leq Cd^2(\tau^h)m(x_0) \leq Cd^2(\tau^h)|v^H|_{1,\infty,\tau_1^h}^2 \leq C|v^H|_{1,\tau_1^h}^2.$$

Summing  $(I)_3$  over all  $\tau^h \in \Omega(x_l^H, x_r^H)$ , and using (4.2)-(4.3), we obtain

$$(4.4) \quad \sum_{\tau^h \in \Omega(x_l^H, x_r^H)} |\mathcal{I}_h v^H|_{1,\tau^h}^2 \leq C \left( |\tilde{\Theta}_h v^H|_{1,\tau_1^H}^2 + |\tilde{\Theta}_h v^H|_{1,\tau_2^H} + |v^H|_{1,N(\tau_{lr}^H)}^2 \right).$$

Note that  $\tilde{\Theta}_h v^H$  is linear over  $\Omega(x_l^H, x_r^H)$ , uniquely determined by values  $v^H(x_l^H)$ ,  $v^H(x_r^H)$  and  $v^H(x_i^H)$ , thus we derive immediately by direct calculations (cf. Fig. 5) that, with  $w^H = \tilde{\Theta}_h v^H$ :

$$|w^H|_{1,\tau_1^H}^2 \leq C \{ (w^H(x_l^H) - w^H(x_r^H))^2 + (w^H(x_r^H) - w^H(x_1^H))^2 + (w^H(x_1^H) - w^H(x_l^H))^2 \}$$

Using the assumption **(H2)**, we know

$$w^H(x_l^H) = v^H(x_l^H), \quad w^H(x_r^H) = v^H(x_r^H).$$

Combining with the definition of  $\Theta_h$ , the boundedness of  $\theta_i$  and (3.1) yields

$$\begin{aligned} |w^H|_{1,\tau_1^H}^2 &\leq C \{ (v^H(x_l^H) - v^H(x_r^H))^2 + (v^H(x_r^H) - v^H(x_i^H))^2 + (v^H(x_i^H) - v^H(x_l^H))^2 \} \\ &\leq C |v^H|_{1,\tau_{lr}^H}^2. \end{aligned}$$

The same result is true for  $|w^H|_{1,\tau_2^H}^2 \equiv |\tilde{\Theta}_h v^H|_{1,\tau_2^H}^2$ . Thus we obtain from these estimates and (4.4) that

$$\sum_{\tau^h \in \Omega(x_l^H, x_r^H)} |\mathcal{I}_h v^H|_{1,\tau^h}^2 \leq C |v^H|_{1,N(\tau_{lr}^H)}^2.$$

This proves (14).  $\square$

Lemma 4.1 plays a crucial role in the convergence analysis of multilevel Schwarz methods, let us just state the following results for the stability and approximation of the coarse space  $V^H$  to the fine space  $V^h$  which immediately give rise to the convergence and condition number bounds for the two-level additive Schwarz methods [7, 8, 9]. As multilevel additive methods need some more tools, for example, stability of the inverse of the coarse-to-fine interpolant and

construction of a good partition of a fine function over the subspaces of all grid levels, we refer to Chan-Zou [9] for more details.

Let  $\hat{\Omega}$  be an open bounded domain in  $R^2$  which is large enough so that it contains both  $\Omega$  and  $\tilde{\Omega}^H$ , and  $E : H^1(\Omega) \rightarrow H^1(\hat{\Omega})$  be a linear extension operator satisfying

$$Ew|_{\Omega} = w, \quad \|Ew\|_{1,\hat{\Omega}} \leq C\|w\|_{1,\Omega}, \quad \forall w \in H^1(\Omega).$$

See Stein [22] for the existence of such an extension operator. Let  $Q_H : L^2(\tilde{\Omega}^H) \rightarrow \tilde{V}^H$  be Clément's interpolant. We refer to Clément [11] for its definition and Chan-Zou [8] and Chan-Smith-Zou [7] for its use in domain decomposition contexts. Evidently,

$$(Q_H w^H)|_{\Omega^H} \in V^H, \quad \forall w^H \in L^2(\tilde{\Omega}^H).$$

LEMMA 4.2. *Given any interpolation operator  $\mathcal{I}_h$  satisfying Definition 3.1, then for any  $u^h \in V^h$  there exists  $u^H \in V^H$  such that for all  $\tau^H \in T^H$ , we have*

$$(11) \quad \sum_{\substack{\tau^h \cap \tau^H \neq \emptyset \\ \tau^h \subset \tilde{\Omega}^H}} \|u^h - \mathcal{I}_h u^H\|_{0,\tau^h}^2 \leq Cd^2(\tau^H) |Eu^h|_{1,N(\tau^H)}^2,$$

$$(12) \quad \sum_{\substack{\tau^h \cap \tau^H \neq \emptyset \\ \tau^h \subset \tilde{\Omega}^H}} |\mathcal{I}_h u^H|_{1,\tau^h} \leq C |Eu^h|_{1,N(\tau^H)},$$

$$(13) \quad \sum_{\tau^h \in \Omega(x_i^H, x_r^H)} \|u^h - \mathcal{I}_h u^H\|_{0,\tau^h}^2 \leq Cd^2(\tau_r^H) \sum_{\tau^H \in N(\tau_r^H)} |Eu^h|_{1,N(\tau^H)}^2,$$

$$(14) \quad \sum_{\tau^h \in \Omega(x_i^H, x_r^H)} |\mathcal{I}_h u^H|_{1,\tau^h}^2 \leq C \sum_{\tau^H \in N(\tau_r^H)} |Eu^h|_{1,N(\tau^H)}^2.$$

*Proof.* As stated in the proof of Lemma 4.1, the proof of the inequalities (11) and (12) is easy and can be found in [7, 8]. We next prove (13) and (14).

For any  $u^h \in V^h$ , we choose  $u^H \in V^H$  by

$$u^H = Q_H Eu^h|_{\Omega^H} \in V^H.$$

This  $u^H$  satisfies the required results. The  $H^1$ -stability (14) is an immediate consequence of Lemma 4.1 and the  $H^1$ -stability of  $Q_H$ . We now prove (13).

On the fine domain  $\Omega^h = \Omega$ , we can split  $u^h - \mathcal{I}_h u^H$  into two parts:

$$(4.5) \quad u^h - \mathcal{I}_h u^H = (Eu^h - Q_H Eu^h) + (Q_H Eu^h - \mathcal{I}_h Q_H Eu^h).$$

*First term estimate in (4.5):* if Neumann boundary condition is imposed at least at one of the two coarse nodes  $x_i^H$  and  $x_r^H$  in the space  $V^H$ , we derive by assumption on  $\Omega(x_i^H, x_r^H)$  and properties of Clément's interpolant  $Q_H$  that

$$\begin{aligned} \sum_{\tau^h \in \Omega(x_i^H, x_r^H)} \|Eu^h - Q_H Eu^h\|_{0,\tau^h}^2 &\leq \|Eu^h - Q_H Eu^h\|_{0,N(\tau_r^H)}^2 \\ &\leq Cd^2(\tau_r^H) \sum_{\tau^H \in N(\tau_r^H)} |Eu^h|_{1,N(\tau^H)}^2. \end{aligned}$$

If Dirichlet boundary condition is imposed on both nodes  $x_i^H$  and  $x_r^H$  in the space  $V^H$ , the result follows from Poincaré inequality.

Second term estimate in (4.5): we obtain from Lemma 4.1 that, with  $v^h = Eu^h$ ,

$$\sum_{\tau^h \in \Omega(x_1^H, x_r^H)} \|Q_H v^h - \mathcal{I}_h Q_H v^h\|_{0, \tau^h}^2 \leq C d^2(\tau_{lr}^H) |Q_H v^h|_{1, N(\tau_{lr}^H)}^2.$$

Then using the stability of  $Q_H$  yields

$$\sum_{\tau^h \in \Omega(x_1^H, x_r^H)} \|Q_H E u^h - \mathcal{I}_h Q_H E u^h\|_{0, \tau^h}^2 \leq C d^2(\tau_{lr}^H) \sum_{\tau^H \in N(\tau_{lr}^H)} |E u^h|_{1, N(\tau^H)}^2.$$

Now (13) follows from (4.5) and the above two estimates for the 1st and 2nd terms in (4.5).  $\square$

**5. Numerical results.** In this section, we provide some numerical results of domain decomposition and multigrid methods on unstructured meshes for elliptic problems on various fine grid domains (see Figure 6). The well-known NASA airfoil mesh was provided by T. Barth and D. Jespersen of NASA Ames, and a fine, unstructured square and annulus were generated using Barth's 2-dimensional Delaunay triangulator. All numerical experiments were performed using the Portable, Extensible Toolkit for Scientific Computation (PETSc) [13], running on a Sun SPARC 20. Piecewise linear finite elements were used for the discretizations and the resulting linear system was solved using either multilevel overlapping Schwarz or V-cycle multigrid as a preconditioner with full GMRES as an outer accelerator.

Our approach to generating a coarse grid hierarchy is to find a maximal independent set of the boundaries and the interior of the fine grid of the mesh, and then retriangulate the resulting set of vertices (other coarsening algorithms can be used here). This process is then repeated recursively for the desired number of levels. An example coarse grid hierarchy of the airfoil mesh retriangulated with Cavendish's algorithm [4] is shown in Figure 7 where  $G^2$  refers to the first coarsening of the fine grid,  $G^1$  is the coarsening of  $G^2$ , and  $G^0$  is the coarsening of the  $G^1$ .

The interpolation matrices are formed by taking each fine node and searching for the coarse grid element in which it lies, then interpolating with the values of the coarse nodes which made up that element using linear interpolation. If the coarse domain does not cover the fine domain, then the linear interpolant,  $\mathcal{I}_h^0$ , leads to a natural extension by zero, which is only appropriate for Dirichlet boundary conditions. If no such coarse grid element can be found (when  $x_j^h \in \Omega \setminus \Omega^H$ ), and a Neumann boundary condition is imposed at the fine node, then one of the interpolants defined as in Section 3 is used ( $\mathcal{I}_h^1$ ,  $\mathcal{I}_h^2$ , or  $\mathcal{I}_h^3$ ). The restriction matrices are taken to be the transposes of the interpolation matrices.




FIG. 6. Some fine grids: an unstructured square with 385 nodes (left), NASA airfoil with 4253 nodes (center) and an annulus with 610 nodes (right).




FIG. 7. Airfoil grid hierarchy with unmodified boundaries (left) and modified boundaries (right).

We shall present numerical results for Schwarz solvers and multigrid methods. For partitioning, all the domains (except the coarsest) were partitioned using the recursive spectral bisection method [20], with exact solves for both the subdomain problems and the coarse grid problem. To generate overlapping subdomains, we first partition the domain into non-overlapping subdomains and then extend each subdomain by some number of elements.

In all the experiments, the initial iterate is set to be zero and the iteration is stopped when the discrete norm of the residual is reduced by a factor of  $10^{-5}$ .

For our first experiment, we use additive Schwarz to solve the Poisson problem on a unit square with homogeneous Dirichlet boundary conditions. Because the fine domain is so simple and Dirichlet boundary conditions are given, non-matching boundaries are not an issue here and no special interpolants are used. We provide these results simply for completeness, as multilevel Schwarz results on unstructured grids have not been previously found in the literature to the authors' knowledge. Table 1 shows the number of GMRES iterations to convergence with varying fine grid problem and varying number of levels. Providing a coarse grid greatly improved convergence, and without it the method is not scalable to larger problems. Interesting things to notice are that for a fixed number of levels, multilevel Schwarz is mesh-size independent, but that the number of iterations increases with the number of levels for a fixed problem size. This had also been previously observed for structured meshes using a multilevel diagonal scaling method in [21] and is due to the additive nature of the method. Also, increasing the amount of overlap improved convergence, but in practice, a one-element overlap was sufficient.

In our second experiment, we solve a mildly varying coefficient problem on the airfoil:

$$\frac{\partial}{\partial x}((1 + xy)\frac{\partial u}{\partial x}) + \frac{\partial}{\partial y}((\sin(3y))\frac{\partial u}{\partial y}) = (4xy + 2)\sin(3y) + 9x^2 \cos(6y)$$

with either a purely Dirichlet boundary condition or a mixed boundary condition: Dirichlet for  $x \leq 0.2$  and homogeneous Neumann for  $x > 0.2$ . For this problem, the non-homogeneous Dirichlet condition is  $u = 2 + x^2 \sin(3y)$ . Table 2 shows the number of GMRES iterations to convergence using additive multilevel Schwarz with the different boundary treatments. We see the slow increase in iteration number as we increase the number of levels used. More importantly, we see the deterioration in the method when Neumann conditions are not properly handled.

In Table 3, we show results for the same problem, but solved using a hybrid multiplicative-additive Schwarz (multiplicative between levels but additive among subdomains on the same level). As in the additive case, deterioration of the method occurs when mixed boundary conditions are present. However, we can achieve optimal convergence rates, even with a varying number of levels with the hybrid method. Still further improvement can be obtained when using a multiplicative method (both on the subdomains and between levels) and the method behaves much like multigrid (see Tables 4-5). In fact, this is nothing more than multigrid but with a block smoother. A V-cycle multigrid method with pointwise Gauss-Seidel smoothing and 2 pre- and 2 post-smoothings per level was used to produce the results in Table 5.

Table 6 shows some multigrid results for the Poisson equation on an annulus. The forcing function is set to be one and both kinds of boundary conditions were tested. A V-cycle multigrid method with pointwise Gauss-Seidel smoothing and 2 pre- and 2 post-smoothings per level was used. When mixed boundary conditions are present, the deterioration is less pronounced in the multigrid method, but it still exists. It is interesting to note that in our previous multigrid experiments on a quasi-uniform annulus (see [5]), the observed deterioration in the method was much more dramatic than those observed here with the unstructured annulus. We believe that this was due to some extremely poor element aspect ratios on the fine grid in the quasi-uniform case, compounding the effect of the poor approximation on Neumann boundaries.

TABLE 1

*Additive multilevel Schwarz iterations for the Poisson problem on a unit square grid. All grids (except coarsest) were partitioned using RSB. Tables show the number of GMRES iterations to convergence.*

Dirichlet boundary conditions					
# of levels	# of nodes	# of subdomains	# overlap elements		
			0	1	2
1	6409	256	84	63	50
	1522	64	45	36	27
	385	16	26	19	16
2	1522	64	19	16	16
	385	1			
	385	16	19	15	15
	102	1			
	102	4	17	15	15
3	29	1			
	6409	256	28	24	25
	1522	64			
	385	1			
	1522	64	32	25	26
	385	16			
	102	1			
385	16	31	26	26	
4	102	4			
	29	1			
	6409	256	43	37	37
	1522	64			
	385	16			
	102	1			
	1522	64	42	37	37
385	16				
102	4				
29	1				

**6. Conclusions.** When using general unstructured meshes, the coarse grid domain may not necessarily match that of the fine grid. For the parts of the fine grid domain which are not contained in the coarse domain, special treatments must be done to handle different boundary conditions. The transfer operators using linear interpolation with a zero extension is the most natural to implement and is effective for problems with Dirichlet boundary conditions.

For problems where Neumann boundary conditions exist however, zero extension is no longer appropriate and special interpolants should be sought. Our numerical results show the significance of the assumption that when standard interpolations with zero extension are used, the coarse grid must cover the Neumann boundaries of the fine grid problem, otherwise deterioration of the methods occurs. The deterioration is most significant when using additive multilevel methods, but can still be seen for the multiplicative methods. When coupled with highly stretched elements, the deterioration can be very significant, even for multiplicative methods.

Although modifying the coarse grid domains to ensure that this assumption is satisfied is

TABLE 2

*Additive multilevel Schwarz iterations for the elliptic problem with mildly varying coefficients on the airfoil grid ( $G^3$ ) with 4253 unknowns. All grids (except coarsest) were partitioned using RSB, with 1 element overlap. Tables show the number of GMRES iterations to convergence. \* indicates identical results since no coarse grid was used.*

Dirichlet boundary conditions						
# of levels	Grids	# of subdomains	Special Interpolant Used			
			$\mathcal{I}_h^0$	$\mathcal{I}_h^1$	$\mathcal{I}_h^2$	$\mathcal{I}_h^3$
1	$G^3$	32	23	*	*	*
2	$G^3$	32	15	15	15	16
	$G^2$	1				
3	$G^3$	32	23	23	23	25
	$G^2$	8				
	$G^1$	1				
4	$G^3$	32	32	33	33	35
	$G^2$	8				
	$G^1$	2				
	$G^0$	1				

Mixed Dirichlet/Neumann boundary conditions						
# of levels	Grids	# of subdomains	Special Interpolant Used			
			$\mathcal{I}_h^0$	$\mathcal{I}_h^1$	$\mathcal{I}_h^2$	$\mathcal{I}_h^3$
1	$G^3$	32	51	*	*	*
2	$G^3$	32	43	14	15	16
	$G^2$	1				
3	$G^3$	32	53	21	23	23
	$G^2$	8				
	$G^1$	1				
4	$G^3$	32	61	27	29	30
	$G^2$	8				
	$G^1$	2				
	$G^0$	1				

effective, this approach can be problematic to implement for particularly complicated domains or can sometimes generate coarse grid domains which deviate significantly from the fine domain.

An alternative is to modify the interpolants so that non-zero extensions be used on those fine grid boundaries which have Neumann conditions and which are not contained within the coarse grid domain. Since we are using the multilevel methods only as preconditioners, the extension need not be particularly accurate; we used either constant extension with the nearest boundary nodal value or extension using the barycentric functions of the nearest coarse grid element, neither of which are difficult to implement. We showed that these more accurate interpolants possess the local  $L^2$ -approximation and  $H^1$ -stability which are essential for proving optimal convergence rates for the multilevel Schwarz methods on unstructured meshes.

## REFERENCES

- [1] R. BANK AND J. XU, *An algorithm for coarsening unstructured meshes*, Numer. Math., 73 (1996), pp. 1-23.

TABLE 3

*Hybrid multiplicative-additive multilevel Schwarz iterations for the elliptic problem with mildly varying coefficients on the airfoil grid ( $G^3$ ) with 4253 unknowns. All grids (except coarsest) were partitioned using RSB, with 1 element overlap. Tables show the number of GMRES iterations to convergence. \* indicates identical results since no coarse grid was used.*

## Dirichlet boundary conditions

# of levels	Grids	# of subdomains	Special Interpolant Used			
			$\mathcal{I}_h^0$	$\mathcal{I}_h^1$	$\mathcal{I}_h^2$	$\mathcal{I}_h^3$
1	$G^3$	32	23	*	*	*
2	$G^3$	32	13	13	13	14
	$G^2$	1				
3	$G^3$	32	13	13	13	14
	$G^2$	8				
	$G^1$	1				
4	$G^3$	32	13	13	13	14
	$G^2$	8				
	$G^1$	2				
	$G^0$	1				

## Mixed Dirichlet/Neumann boundary conditions

# of levels	Grids	# of subdomains	Special Interpolant Used			
			$\mathcal{I}_h^0$	$\mathcal{I}_h^1$	$\mathcal{I}_h^2$	$\mathcal{I}_h^3$
1	$G^3$	32	51	*	*	*
2	$G^3$	32	36	13	13	14
	$G^2$	1				
3	$G^3$	32	36	13	13	14
	$G^2$	8				
	$G^1$	1				
4	$G^3$	32	36	13	13	14
	$G^2$	8				
	$G^1$	2				
	$G^0$	1				



TABLE 4

*Multiplicative multilevel Schwarz iterations for the elliptic problem with mildly varying coefficients on the airfoil grid ( $G^3$ ) with 4253 unknowns. All grids (except coarsest) were partitioned using RSB, with 1 element overlap. Tables show the number of GMRES iterations to convergence. \* indicates identical results since no coarse grid was used.*

Dirichlet boundary conditions						
# of levels	Grids	# of subdomains	Special Interpolant Used			
			$\mathcal{I}_h^0$	$\mathcal{I}_h^1$	$\mathcal{I}_h^2$	$\mathcal{I}_h^3$
1	$G^3$	32	9	*	*	*
2	$G^3$	32	4	4	4	4
	$G^2$	1				
3	$G^3$	32	4	4	4	4
	$G^2$	8				
	$G^1$	1				
4	$G^3$	32	4	4	4	4
	$G^2$	8				
	$G^1$	2				
	$G^0$	1				

Mixed Dirichlet/Neumann boundary conditions						
# of levels	Grids	# of subdomains	Special Interpolant Used			
			$\mathcal{I}_h^0$	$\mathcal{I}_h^1$	$\mathcal{I}_h^2$	$\mathcal{I}_h^3$
1	$G^3$	32	23	*	*	*
2	$G^3$	32	5	4	4	4
	$G^2$	1				
3	$G^3$	32	5	4	4	4
	$G^2$	8				
	$G^1$	1				
4	$G^3$	32	5	4	4	4
	$G^2$	8				
	$G^1$	2				
	$G^0$	1				

- [2] J. BRAMBLE, J. PASCIAK, AND J. XU, *Parallel multilevel preconditioners*, Math. Comp., 55 (1990), pp. 1–21.
- [3] X.-C. CAI AND Y. SAAD, *Overlapping domain decomposition algorithms for general sparse matrices*, Numerical Linear Algebra with Applications, 3 (1996), pp. 221–237.
- [4] J. C. CAVENDISH, *Automatic triangulation of arbitrary planar domains for the finite element method*, Int'l J. Numer. Meth. Engineering, 8 (1974), pp. 679–696.
- [5] T. F. CHAN, S. GO, AND J. ZOU, *Multilevel domain decomposition and multigrid methods for unstructured meshes: Algorithms and theory*, Tech. Rep. 95–24, Department of Mathematics, University of California at Los Angeles, May 1995. also in Proceedings of the Eighth International Conference on Domain Decomposition, May 1995, Beijing.
- [6] T. F. CHAN AND B. SMITH, *Domain decomposition and multigrid methods for elliptic problems on unstructured meshes*, in Domain Decomposition Methods in Science and Engineering, Proceedings of the Seventh International Conference on Domain Decomposition, October 27–30, 1993, The Pennsylvania State University, D. Keyes and J. Xu, eds., American Mathematical Society, Providence, 1994. also in Electronic Transactions

TABLE 5

*Multigrid iterations for the elliptic problem with mildly varying coefficients on the airfoil. Tables show the number of GMRES iterations to convergence.*

Dirichlet boundary conditions						
# of fine grid nodes	MG levels	# of coarse grid nodes	Special Interpolant Used			
			$\mathcal{I}_h^0$	$\mathcal{I}_h^1$	$\mathcal{I}_h^2$	$\mathcal{I}_h^3$
4253	2	1170	4	4	4	4
	3	340	4	4	4	4
	4	101	4	4	4	4

Mixed Dirichlet/Neumann boundary conditions						
# of fine grid nodes	MG levels	# of coarse grid nodes	Special Interpolant Used			
			$\mathcal{I}_h^0$	$\mathcal{I}_h^1$	$\mathcal{I}_h^2$	$\mathcal{I}_h^3$
4253	2	1170	6	5	4	4
	3	340	6	4	5	5
	4	101	7	5	5	5

TABLE 6

*Multigrid iterations for the Poisson problem on an annulus. The exit condition was decreased to  $10^{-6}$  from  $10^{-5}$ . Tables show the number of GMRES iterations to convergence.*

Dirichlet boundary conditions						
# of fine grid nodes	MG levels	# of coarse grid nodes	Special Interpolant Used			
			$\mathcal{I}_h^0$	$\mathcal{I}_h^1$	$\mathcal{I}_h^2$	$\mathcal{I}_h^3$
2430	2	610	4	4	4	4
	3	160	4	4	4	4
	4	47	4	4	4	4

Mixed Dirichlet/Neumann boundary conditions						
# of fine grid nodes	MG levels	# of coarse grid nodes	Special Interpolant Used			
			$\mathcal{I}_h^0$	$\mathcal{I}_h^1$	$\mathcal{I}_h^2$	$\mathcal{I}_h^3$
2430	2	610	6	5	4	4
	3	160	7	5	4	4
	4	47	7	5	4	4

- on Numerical Analysis, v.2, (1994), pp. 171–182.
- [7] T. F. CHAN, B. SMITH, AND J. ZOU, *Overlapping Schwarz methods on unstructured meshes using non-matching coarse grids*, Numerische Mathematik, 73 (1996), pp. 149–167.
- [8] T. F. CHAN AND J. ZOU, *Additive Schwarz domain decomposition methods for elliptic problems on unstructured meshes*, Numerical Algorithms, 8 (1994), pp. 329–346.
- [9] —, *A convergence theory of multilevel additive Schwarz methods on unstructured meshes*, Tech. Rep. 95-16, Department of Mathematics, University of California at Los Angeles, March 1995. to appear in Numerical Algorithms.
- [10] P. CIARLET, *The Finite Element Method for Elliptic Problems*, North-Holland, 1978.
- [11] P. CLÉMENT, *Approximation by finite element functions using local regularization*, R.A.I.R.O. Numer. Anal., R-2 (1975), pp. 77–84.
- [12] H. DECONNICK AND T. BARTH, eds., *Special course on unstructured grid methods for advection dominated flows*, March 1992. AGARD REPORT 787, Special course at the

- VKI, Belgium.
- [13] W. D. GROPP AND B. F. SMITH, *Portable Extensible Toolkit for Scientific Computation (PETSc)*. Available by anonymous ftp at [info.mcs.anl.gov](ftp://info.mcs.anl.gov) in the directory `pub/pdetools` or at <http://www.mcs.anl.gov/Projects/petsc/petsc.html>.
  - [14] H. GUILLARD, *Node-nested multi-grid method with Delaunay coarsening*, Tech. Rep. RR-1898, INRIA, Sophia Antipolis, France, March 1993.
  - [15] B. KOOBUS, M. H. LALLEMAND, AND A. DERVIEUX, *Unstructured volume-agglomeration MG: solution of the Poisson equation*, International Journal for Numerical Methods in Fluids, 18 (1994), pp. 27–42.
  - [16] R. KORNUBER AND H. YSERENTANT, *Multilevel methods for elliptic problems on domains not resolved by the coarse grid*, in Domain Decomposition Methods in Science and Engineering, Proceedings of the Seventh International Conference on Domain Decomposition, October 27–30, 1993, The Pennsylvania State University, D. Keyes and J. Xu, eds., vol. 180, American Mathematical Society, Providence, 1994, pp. 49–60.
  - [17] D. MAVRIPLIS, *Unstructured mesh algorithms for aerodynamic calculations*, Tech. Rep. 92-35, ICASE, NASA Langley, Virginia, July 1992.
  - [18] D. J. MAVRIPLIS AND V. VENKATAKRISHNAN, *Agglomeration multigrid for two-dimensional viscous flows*, Computers & Fluids, 24 (1995), pp. 553–570.
  - [19] C. F. OLLIVIER-GOOCH, *Multigrid acceleration of an upwind Euler solver on unstructured meshes*, AIAA Journal, 33 (1995), pp. 1822–1827.
  - [20] A. POTHEN, H. SIMON, AND K. LIU, *Partitioning sparse matrices with eigenvector of graphs*, SIAM J. Mat. Anal. Appl., 11 (1990), pp. 430–452.
  - [21] B. SMITH, P. BJØRSTAD, AND W. GROPP, *Domain Decomposition*, Cambridge University Press, Cambridge, 1996.
  - [22] E. M. STEIN, *Singular integrals and differentiability properties of functions*, Princeton University Press, Princeton, 1970.
  - [23] J. XU, *Iterative methods by space decomposition and subspace correction*, SIAM Review, 34 (1992), pp. 581–613.
  - [24] X. ZHANG, *Multilevel Schwarz methods*, Numer. Math., 63 (1992), pp. 521–539.