

Basic Exam: Fall 2024

Test instructions:

- Write your UCLA ID number on the upper right corner of each page.
- Do not write your name anywhere on the exam.
- Your final score will be the sum of
FIVE linear algebra problems (Problems 1-6) and
FIVE analysis problems (Problems 7-12).
However, to pass the exam you need to show mastery of both subjects.
- Indicate below which 10 problems you wish to have graded.
- Please staple your problems in numerical order.

1	2	3	4	5	6
7	8	9	10	11	12

Linear Algebra Problems

Problem 1. Let $A = \{a_{ij}\}_{i,j=1}^d$ be an integer-valued matrix whose row sums are all equal to a positive integer n . Noting that $\det(A)$ is an integer, prove that n divides $\det(A)$.

Problem 2. Let $n \geq 2$ be an integer and A be a normal complex-valued $n \times n$ -matrix. Assume that $\dim(\text{Ker}(\lambda - A)) \leq 1$ for all $\lambda \in \mathbb{C}$. Prove that any normal complex-valued $n \times n$ -matrix B that commutes with A takes the form

$$B = b_0 \text{Id} + b_1 A + \cdots + b_{n-1} A^{n-1}$$

for some $b_0, \dots, b_{n-1} \in \mathbb{C}$.

Problem 3. Suppose that T is a linear operator on a finite-dimensional complex vector space with spectrum $\sigma(T) = \{0\}$. Show that T is nilpotent.

Problem 4. Consider the polynomial

$$p(x) = x^4 + 2x^3 - 3x^2 - 4x + 4$$

(To save you some time, note that this polynomial vanishes only at $x = 1$ and $x = -2$.) Let A be a complex-valued 4×4 -matrix such that $p(A) = 0$. Assume also that $\text{Tr}(A) = 1$ and $\text{Rank}(A - \text{Id}) = 2$. Provide the Jordan canonical form for A . Justify your answer.

Problem 5. Solve the system of ODEs for functions $x(t)$ and $y(t)$:

$$\begin{aligned}x'(t) &= 2x(t) + y(t) \\y'(t) &= x(t) - 2y(t)\end{aligned}$$

with initial data $x(0) = 1$ and $y(0) = 0$.

Problem 6. Let V be a linear vector space of functions on a set X of finite cardinality $|X|$. Prove that $\dim(V) \leq |X|$. Then prove existence of points $x_1, \dots, x_n \in X$, with $n = \dim(V)$, and functions $f_1, \dots, f_n \in V$ such that

$$f_i(x_j) = \delta_{ij}, \quad i, j = 1, \dots, n$$

(In particular, f_1, \dots, f_n is a basis in V .)

Analysis Problems

Problem 7. Prove that given any continuous, increasing function $f: \mathbb{R} \rightarrow \mathbb{R}$ with

$$\lim_{x \rightarrow \infty} f(x) = \infty,$$

there exists a differentiable, strictly increasing function $g: \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\lim_{x \rightarrow \infty} g(x) = \infty, \quad \lim_{x \rightarrow \infty} \frac{g(x)}{f(x)} = 0.$$

Problem 8. Let $a < b$ be reals. Prove that if $f: [a, b] \rightarrow \mathbb{R}$ is a continuous and one-to-one function, then f is monotone.

Problem 9. Let $\mathbb{N} = \{1, 2, 3, \dots\}$. Consider two real-valued sequences $\{a_n\}_{n \in \mathbb{N}}$ and $\{b_n\}_{n \in \mathbb{N}}$ which satisfy the following two properties:

(i) There exists a constant $A > 0$ such that for all $n \in \mathbb{N}$ the following holds:

$$\left| \sum_{i=1}^n a_i \right| \leq A$$

(ii) The sequence $\{b_n\}_{n \in \mathbb{N}}$ is monotone decreasing and $\lim_{n \rightarrow \infty} b_n = 0$.

Do the following:

(a) Show that, for all $m \in \mathbb{N}$,

$$\sum_{i=n}^m a_i b_i = F_m b_m - F_{n-1} b_n + \sum_{i=n}^{m-1} F_i (b_i - b_{i+1})$$

holds with $F_k := \sum_{i=1}^k a_i$.

(b) Use part (a) to prove that the series $\sum_{n=1}^{\infty} a_n b_n$ converges.

Problem 10. Let X and Y be metric spaces, with distance functions d_X and d_Y , respectively. Prove that if X is compact, then any continuous function $f: X \rightarrow Y$ is uniformly continuous. For full credit, solve this problem directly from the relevant definitions, without reference to any major theorems.

Problem 11. (a) Suppose $F(x, y)$ is a continuous function on \mathbb{R}^2 such that for every rectangular domain $D = [a, b] \times [c, d]$ the following holds:

$$\iint_D F(x, y) dx dy = 0.$$

Prove that $F(x, y) = 0$ for all $(x, y) \in \mathbb{R}^2$.

(b) Assume that $f(x, y)$, $\frac{\partial}{\partial x} f(x, y)$, $\frac{\partial}{\partial y} f(x, y)$, $\frac{\partial}{\partial x} \left(\frac{\partial}{\partial y} f(x, y) \right)$, $\frac{\partial}{\partial y} \left(\frac{\partial}{\partial x} f(x, y) \right)$ are all continuous on \mathbb{R}^2 . Use part (a) of this problem to prove that:

$$\frac{\partial}{\partial y} \left(\frac{\partial}{\partial x} f(x, y) \right) = \frac{\partial}{\partial x} \left(\frac{\partial}{\partial y} f(x, y) \right).$$

You may assume without proof that the iterated integrals $\iint F(x, y) dx dy$ and $\iint F(x, y) dy dx$ are equal.

Problem 12. Consider a sequence $\{f_n\}_{n \in \mathbb{N}}$ of functions $f_n: \mathbb{R} \rightarrow \mathbb{R}$ such that

- (i) f_n has at most one discontinuity point for each $n \in \mathbb{N}$, and
- (ii) $f_n \rightarrow f$ uniformly for some function $f: \mathbb{R} \rightarrow \mathbb{R}$.

Prove that f has at most one discontinuity point.